

MARCH 1, 1950

# MODERN MEDICINE

*The Journal of Diagnosis and Treatment*



Dr. R. A. Gordon  
(see page 9)

*Isidor Krollman*

Table of contents  
page 9



## The most "persuasive" oral germicide you can prescribe

1. Cēpacol persuades a wide range of oral bacteria to surrender within 15 seconds after contact<sup>1</sup>
2. Cēpacol's pleasant taste persuades your patients to use it

The rapid antiseptics<sup>2</sup> and soothing relief which Cēpacol brings to inflamed, sore throats are important. Along with the fact that Cēpacol is non-irritating, non-toxic, and does not interfere with tissue healing. Too, patients are extremely grateful to you for prescribing something so effective that also is so pleasant to use—as either gargle or spray.

# CĒPACOL®

The alkaline germicidal solution that works in partnership with saliva



CINCINNATI • U.S.A.

**NOW AVAILABLE**—Cēpacol Throat Lozenges! These convenient, pleasant-tasting lozenges, dissolved slowly in the mouth, provide a soothing, analgesic solution to relieve the dryness and irritation of sore throat.


1. As shown in laboratory studies. 2. Cēpacol contains an effective germicidal detergent, the quaternary ammonium salt Cetylpyridinium Chloride, 1:5000.

# 20-minute diagnosis of pregnancy

WITH THE

*Westerfield®*

## CHEMICAL PREGNANCY TEST\* (Carson-Sacks Method)



### 94.4% accurate in 431 cases reported<sup>1</sup>

This recent report<sup>1</sup> on a large series of unselected cases not only confirms the reliability of the Westerfield Chemical Pregnancy Test in the early diagnosis of pregnancy, but also emphasizes its simplicity.

#### FEATURES:

- New, simple office procedure—only apparatus required is a beaker and hot plate.
- A urine test—one or several tests can be run in twenty to thirty minutes.
- No delay—results can be read immediately.
- Economical—costs less per test than other reliable methods.

SUPPLIED: 12-test, 45-test, and 225-test sets in attractive compact cases; refill units; individual replacement parts.

Detailed Literature Available.

\*U. S. Pat. Applied For.

1. Ricketts, W. A.; Carson, R. M., and Sacks, R. R.: Am. J. Obst. & Gynec. 56:955 (Nov.) 1948.

**BEUCHLER'S**  
**COLUMBIA MEDICAL BUILDING ANNEX**  
Republic 6007

915 19th STREET, N. W.

WASHINGTON, D. C.

*You've asked for it...*



**RECEPTION ROOM FURNITURE**

- **DESKS**
- **SIDE CHAIRS**
- **ARM CHAIRS**
- **DAVENPORTS**
- **SETTEES**
- **END TABLES**
- **MAGAZINE TABLES**

You've asked for it ... and Hamilton provides the answer in this beautiful new reception room and business office furniture. Here is furniture that lends a competent and professional air to your office. At the same time, it provides the comforts your patients expect. Sturdy, No-Sag construction, comfortable padding. Rich, durable, easy-to-clean plastic covering in a wide selection of inviting colors. See the complete line today!

**BEUCHLER'S**

**COLUMBIA MEDICAL BUILDING ANNEX**

Republic 6007

915 19th STREET, N. W.

WASHINGTON, D. C.



**20% BENZOCAINE IN SOLUTION**

# *Americaine*

**TOPICAL ANESTHETIC OINTMENT**

Clear or with Chlorophyll

**MOST POTENT.** 20% Benzocaine<sup>1</sup> is NOW dissolved in a bland, water-soluble ointment—heretofore only 2% was generally considered possible. Furthermore, to be effective, Benzocaine must be in solution.

**TAINTER<sup>2</sup>**, in a series of studies involving about 40 different materials, found Benzocaine to be "the best topical anesthetic."

**LEAST TOXIC.** While Benzocaine's action is generally recognized for its lack of toxicity, **ADRIANI<sup>3</sup>**, in evaluating the relative toxicity of various topical anesthetics, found Benzocaine to be by far the safest.

**PROLONGED ANESTHESIA.** Controlled clinical studies reveal 4-6 hours' relief from pruritus associated with dermatoses, exanthemas, diabetes, etc.

Americaine Clear (Benzocaine 20%, oxyquinoline benzoate, 0.39%); with Chlorophyll (same constituents with 0.08% chlorophyll added to stimulate cellular regeneration).



- References. 1. Finkel, M., Levine, A. J., Wohl, M. Twenty percent dissolved benzocaine ointment in the treatment of burns. *Ind. Med.* 17: 475 Dec., 1948.  
2. Tainter, M. L. Some general considerations in evaluating local anesthetic solutions in patients. *Anesthesiology* 3: 470 Sept. 1944.  
3. Adriani, J. The pharmacology of anesthetic drugs. *Chas. C. Thomas*, 1941, p. 49.

# *Americaine*

... is available in two forms: Clear and with Chlorophyll. Supplied in 1 ounce tubes or ½ lb., 1 lb. and 7 lb. jars.

**BEUCHLER'S**  
**COLUMBIA MEDICAL BUILDING ANNEX**  
Republic 6007

915 19th STREET, N. W.

WASHINGTON, D. C.

**NEW!**... and more effective for

- COAGULATION
- DESICCATION
- FULGURATION

**The NATIONAL  
ELECTRICATOR \***

Always ready for instant use. Simply pick up handle—press the button—your *Electricator*\* goes to work. No electrical connections to worry about . . . no fumbling for foot switch.

One handle takes both bi-polar and mono-polar applicator needles interchangeably. Handle and needles conveniently in place always. Power intensity regulated by "attenuator-control" knob — permits spark gaps to be factory pre-set for optimum Faradic-free efficiency.

A complete instrument self-contained and convenient to mount on wall, occupying less than half the space of ordinary letterhead. Lustrous walnut housing . . . handsome ivory handle and knobs.

**\$45<sup>50</sup>** Choice of any  
2 of 6 Electrodes



**NO FOOT SWITCH FUMBLING!**

Simplicity of design makes *NATIONAL'S* new *ELECTRICATOR*® outstanding for functional efficiency. Easy, fast, comfortable to use — assuring successful results.

\*Trade Mark

Complete with brochure, "Digest of Treatment Procedures."

**BEUCHLER'S**  
**COLUMBIA MEDICAL BUILDING ANNEX**  
Republic 6007

915 19th STREET, N. W.

WASHINGTON, D. C.

IN DAYTIME SEDATION



MENOPAUSAL HYSTERIA



HYPERTENSION



OBSTETRICAL HYPNOSIS



PREOPERATIVE APPREHENSION



INSOMNIA



# butisol sodium

BRAND OF BUTABARBITAL SODIUM

*intermediate sedative*

Butisol Sodium produces "a relatively mild and more continuous depression than can be obtained with the shorter-acting barbiturates, yet its action is less prolonged than with barbital or phenobarbital."

With proper regulation of dosage, there is no cumulative action and a minimum of lethargy and "hang-over."

Sedation is sustained for approximately five to six hours—without sharp peaks of effect—thus providing a most useful sedative-hypnotic in a wide range of clinical indications.

**DOSAGE FORMS:** Elixir Butisol Sodium 0.2 Gm. (3 gr.) per fl. oz. Also Capsules and Tablets, 0.1 Gm. (1½ gr.); Tablets, 15 mg. (¼ gr.), 30 mg. (½ gr.) and 50 mg. (¾ gr.).

*Elixir*  
**butisol  
sodium**

Its bright, green color and refreshing flavor appeal to all; excellent prescription vehicle. Clinical samples on request.



1. J.A.M.A. 135:224 (Sept. 27) 1947.

**McNEIL**


LABORATORIES, INC., PHILADELPHIA 32, PA.

# NOVALENE

Reg. U. S. Pat. Off.

**NOVALENE—Decongestive Tablets—**A time-tested, dependable therapeutic aid provides rapid relief of distressing symptoms in asthma and hay fever without inducing drowsiness or other undesirable side-effects.

**SUPPLIED:** In packages of 25 and 100 tablets, on prescription. Available through all reliable pharmacies. Literature and samples to physicians on request.



**PROFESSIONAL DRUGS, INC.**

*Sole Distributors*

76 NINTH AVE., NEW YORK 11, N. Y.

COUNCIL ACCEPTED

## Cellothyl

### THE CONSTIPATION CORRECTIVE WHICH

Bargen found relieved "obstinate constipation of long duration"  
—often in one to two weeks.<sup>1</sup>

Schweig reported achieved "excellent" or "good" results in  
92% of acute and chronic cases of 1 week to 15 years'  
duration.<sup>2</sup>

### A NEW HYDROPHILIC COLLOID

declared by the J. A. M. A. to have "obvious advantages." It is  
"nontoxic... not absorbed by the intestinal mucosa...  
not degraded by intestinal enzymes... nonantigenic."<sup>3</sup>

shown by Tainter to increase "the bulk of the stools about 10 g."  
for each gram ingested.<sup>4</sup>



## Cellothyl

CHILCOTT  
*Laboratories*

The Maltine Company

# MODERN MEDICINE

THE JOURNAL OF DIAGNOSIS AND TREATMENT

## EDITORIAL BOARD

James T. Case, M.D., *Chicago*  
Franklin D. Dickson, M.D., *Kansas City*  
Julius Hess, M.D., *Chicago*  
Walter B. Hoover, M.D., *Boston*  
Foster Kennedy, M.D., *New York City*  
John C. Krantz, Jr., Ph.D., *Baltimore*  
A. J. Lanza, M.D., *New York City*  
Milton S. Lewis, M.D., *Nashville*  
George R. Livermore, M.D., *Memphis*  
Francis W. Lynch, M.D., *St. Paul*  
Cyril M. MacBryde, M.D., *St. Louis*  
Karl A. Meyer, M.D., *Chicago*  
Alton Ochsner, M.D., *New Orleans*  
Robert F. Patterson, M.D., *Knoxville*  
Edwin B. Plimpton, M.D., *Los Angeles*  
Fred W. Rankin, M.D., *Lexington, Ky.*  
John Alton Reed, M.D., *Washington*  
Rufus S. Reeves, M.D., *Philadelphia*  
Dalton K. Rose, M.D., *St. Louis*  
Howard A. Rusk, M.D., *New York City*  
Roger S. Siddall, M.D., *Detroit*  
James S. Simmons, M.D., *Boston*  
W. Calhoun Stirling, M.D., *Washington*  
Frank P. Strickler, M.D., *Louisville*  
Richard Torpin, M.D., *Augusta, Ga.*  
Paul M. Wood, M.D., *New York City*

---

SCIENCE WRITERS: Thomas B. Gibbons, M.D.,  
Jorma M. Leinassar, M.D., Norman  
Shrifter, M.D., Bernardine Lufkin, Pa-  
tricia O. Read, Elizabeth Kane, Edward  
S. Niederkorn, David W. Morgan

TRANSLATORS: John W. Brennan, M.D.

## EDITORIAL CONSULTANTS

Carolyn G. Adams, M.D.  
E. R. Anderson, M.D.  
Joe W. Baird, M.D.  
Edward P. Burch, M.D.  
James B. Carey, M.D.  
C. D. Creevy, M.D.  
C. J. Ehrenberg, M.D.  
W. K. Haven, M.D.  
Charles Hymes, M.D.  
Miland E. Knapp, M.D.  
Ralph T. Knight, M.D.  
John F. Pohl, M.D.  
Leo Rigler, M.D.  
Wallace P. Ritchie, M.D.  
M. B. Sinykin, M.D.  
A. V. Stoesser, M.D.  
Arthur L. H. Street, LL.B.  
Marvin Sukov, M.D.  
Harry A. Wilmer, M.D.

## EDITORIAL STAFF

A. E. Hedback, M.D.  
*Editor*  
Thomas Ziskin, M.D.  
*Associate Editor*  
Maurice B. Visscher, M.D.  
*Consultant in Sciences*  
Mark S. Parker  
*Executive Editor*  
Sarah A. Davidson  
*Managing Editor*  
Daisy Stilwell  
*Art Editor*  
James Niess  
*Editorial Board  
Secretary*  
Margaret W. Brodt  
Eva Murray  
*Editorial Assistants*

# S.K.F. announces Dexamyl

a balanced combination  
of 'Dexedrine'  
& 'Amytal'

***the remarkable new preparation for  
relieving mental and emotional distress***

*In 'Dexamyl'\*, the two components—'Dexedrine'\* and 'Amytal'†—  
work together to ameliorate mood; to relieve inner tension;  
and thus to control troublesome symptoms  
of mental and emotional distress:*

*The 'Dexedrine', because of its "smooth" and profound  
antidepressant action, restores mental alertness and  
optimism and dispels psychogenic fatigue.*

*The 'Amytal', because of its calming action, relieves  
nervous tension, anxiety and agitation.*

*Widely useful in everyday practice, 'Dexamyl' tablets  
are available on prescription only in bottles of 50.*

*Each tablet contains 'Dexedrine' Sulfate  
(dextro-amphetamine sulfate, S.K.F.) 5 mg. and 'Amytal'  
(Amobarbital, Lilly) ½ grain (32 mg.).*

*\*Trademark, S.K.F. †Trademark, Lilly*

Smith, Kline & French Laboratories, Philadelphia



**The best years  
of their lives . . .**



For almost a decade now they've had Vi-Penta Drops to help them grow. This pioneer water-miscible multivitamin drop preparation protects them in the rapid growth years with a generous supplement of vitamin C and members of the B complex, *in addition to A-and-D*. Vi-Penta Drops are freely miscible with milk and fruit juices. They are easily administered, well-tolerated and well-absorbed. Available in vials of 15 cc, 30 cc, and 60 cc.

HOFFMANN-LA ROCHE INC. • NUTLEY 10 • N. J.

**Vi-Penta® Drops**

**'Roche'**

CORRESPONDENCE.....	14
FORENSIC MEDICINE.....	28
QUESTIONS & ANSWERS.....	38
SPECIAL ARTICLE	
Anesthetic Disasters	
<i>R. A. Gordon</i> .....	45
MEDICINE	
Aureomycin Therapy of Herpes Zoster	
<i>Maxwell Finland, Edmund F. Finnerty,</i>	
<i>Jr., Harvey S. Collins, John W. Baird,</i>	
<i>Thomas M. Gocke, and Edward H. Kass</i>	49
Venous Stasis in the Legs	
<i>W. J. Merle Scott and</i>	
<i>Michael Radakovich</i> .....	50
Guaiac Test for Occult Blood.....	51
Chilblains	
<i>John T. Ingram</i> .....	52
PEDIATRICS	
Diagnosis of Mumps.....	53
Whooping Cough.....	53
Brain Tumors in Children	
<i>A. Earl Walker and Theron L. Hopple</i>	54
Bacterial Pneumonia.....	55
GERIATRICS	
Hyperthyroidism after Sixty	
<i>Elmer C. Bartels and J. W. Kingsley, Jr.</i>	56
GYNECOLOGY & OBSTETRICS	
Functional Sterility and Amenorrhea	
<i>Rita S. Finkler</i> .....	57
Causes and Effects of Eclampsia	
<i>Charles I. Bryans, Jr.,</i>	
<i>and Richard Torpin</i> .....	58

THE MAN ON THE COVER is Dr. R. A. Gordon, general secretary of the Canadian Anaesthetists' Society and author of the Special Article, "Anesthetic Disasters," on page 45 of this issue. Dr. Gordon has written many articles relating to the science of anesthesia and his introduction of the intravenous administration of procaine in treatment of burns attracted world-wide attention. During the war he served overseas as anesthetist in the Royal Canadian Army Medical Corps. He still serves as consulting anesthetist to the Toronto Military Hospital, in addition to carrying on an active practice and teaching at the University of Toronto.



## Contents for March 1 1950

VOL. 18, NO. 5





*Contents  
for  
March 1  
1950*

CONTINUED

**SURGERY**

- A Nonmoving Ligature-Holding Bobbin  
*John Devine*..... 59

**SURGICAL TECHNIGRAM**

- Hemorrhoidectomy  
*F. M. Al Akl*..... 60

**SURGERY**

- Blood Pressure and Surgical Position  
*L. H. Peterson, Kenneth F. Eather,  
and Robert D. Dripps*..... 64

**UROLOGY**

- Serum Acid Phosphatase..... 65  
Bacillary Urinary Infections..... 65

**MEDICAL FORUM**

- Abdominal Paracentesis..... 68  
Temperature Cycle in Obstetrics  
and Gynecology..... 69  
Amyotrophic Lateral Sclerosis..... 69  
Diet in Diabetes..... 70  
Culdoscope in Gynecology..... 72  
What Can Be Done for  
the Hay Fever Patient?..... 73

**DIAGNOSTIX**..... 74

**SHORT REPORTS**..... 80

**MEDICAL NEWS**

- The Accident-prone Automobile Driver  
*W. A. Tillmann and G. E. Hobbs*..... 100

**WASHINGTON LETTER**..... 108

**CURRENT BOOKS & PAMPHLETS**..... 118

**PATIENTS I HAVE MET**..... 124

**MODERN MEDICINE**, The Journal of Medical Progress, is published twice monthly on the first and fifteenth of each month at Minneapolis, Minn. Subscription rate: \$5.00 a year, 25c a copy. Business Manager: M. E. Herz. Address editorial correspondence to 84 South 10th Street, Minneapolis 5, Minn. Telephone: Bridgeport 1291. ADVERTISING REPRESENTATIVES: New York 17: Lee Klemmer, George Doyle, Bernard A. Smiler, 1 East 42nd Street, Suite 801, Corn Exchange Bank Bldg. Telephone: Murray Hill 2-8717. CHICAGO 6: Jay H. Herz, 20 North Wacker Drive, Suite 1921. Telephone: Central 6-4619. SAN FRANCISCO 4: Duncan A. Scott & Co., Mills Bldg. Telephone: Garfield 1-7950. LOS ANGELES 5: Duncan A. Scott & Co., 2978 Wilshire Blvd. Telephone: Dunkirk 8-4151.

**In Eradication and  
Control of TRICHOMONIASIS**  
**ARGYPULVIS is**  
**98% EFFECTIVE\***

\*Reich, Button and Nechtow in  
*Surgery, Gynecology and Obstetrics.*  
Reprint on request.



**THIS** gratifying clinical evidence may be verified in everyday practice with the ARGYPULVIS therapy.

This newer adaptation of ARGYROL lends itself perfectly to the objective: (1) because of its positive bacteriostatic,

detergent and demulcent actions, and (2) because of its convenient forms for office and supplementary home use. Let a professional sample in your office re-demonstrate the reported clinical effectiveness of this product.

.....in two convenient forms.....



**For Use by the Physician.** 7-gram bottles fitting Holm-spray or equivalent powder-blower (in cartons of 3)

**For Home Use by the Patient** 2-gram capsule for insertion by the patient (in bottles of 12)



**INTRODUCTORY TO PHYSICIANS:** \*On request we will send professional samples of ARGYPULVIS (both forms), together with a reprint of the Reich, Button and Nechtow report. (Use coupon.)

A. C. Barnes Company  
Dept. MM-39, New Brunswick, N. J.

Name .....

Address .....

City.....State.....

**ARGYPULVIS**

ARGYROL and ARGYPULVIS are registered trademarks, the property of  
**A. C. BARNES CO., NEW BRUNSWICK, N. J.**



✓ palatable

✓ potent

✓ complete

SYRUP

# Hyotole<sup>®</sup>

*vitamin B-complex and ferrous sulfate*

SHARP & DOHME \_\_\_\_\_

# check list to check ANEMIA

These food factors, as combined in Sharp & Dohme's potent new antianemic preparation, HYOTOLE Syrup, produce an exceptionally prompt hemopoietic response in all types of anemia susceptible to oral treatment:

1 fl. oz. (30 cc.) HYOTOLE Syrup Provides			
✓		✓	
✓		✓	
✓		✓	
✓		✓	
✓		✓	

HYOTOLE Syrup is pleasantly flavored and particularly acceptable to children, as well as to obstetric and geriatric patients. It is especially indicated in nutritional anemias. Supplied in *Spasaver*® pints, and in gallon bottles. Sharp & Dohme, Philadelphia 1, Pa.

# Correspondence

*Communications from the readers of MODERN MEDICINE are always welcome. Address communications to The Editors of MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.*

## **Iodized Oil Safer**

TO THE EDITORS: I always enjoy looking over *Modern Medicine* and in a recent issue (Jan. 1, 1950, p. 92) I was especially interested in the Diagnostix report of the patient with the spontaneous rupture of the esophagus.

There is one correction that I would like to make in Part IV of the discussion and that is concerning the suggestion that a swallow of barium might be used to aid in diagnosis. Barium should not be given in suspected rupture of the esophagus or in patients having complete or almost complete esophageal obstruction. Iodized oil is just as satisfactory and is much safer.

PORTER P. VINSON, M.D.  
Richmond, Va.

## **Intravenous Route During Shock**

TO THE EDITORS: While I was serving in the Army as a surgeon in a station hospital, we developed and used a technic which I would like to pass on to any of my colleagues who may be interested.

In the course of two years we saw many patients who entered our hospital in various stages of primary and secondary shock. As we all know, giving intravenous medications to patients with collapsed venous systems

is a problem. The situation requires a "cut-down" on a vein with the insertion of a catheter or some such procedure.

Use of the corpora cavernosa of the penis has been reported, but I do not think this convenient route has been given the recognition it deserves. We used it many times in plasma and blood transfusions, thereby saving much time, which is so vital in all types of shock.

From our experience, the network of the corpora will take blood or plasma as fast as any vein. For intravenous medication the process is even easier, as a smaller needle may be used. This route was also used with success in male infants who needed transfusions and was much simpler than trying to employ head veins. In all of our cases we had no infections or sequelae of any kind except, perhaps, a mild disapproval of the nursing staff.

The penis may be prepared and made as sterile as possible by any convenient method. We used tincture of green soap and followed up with merthiolate. The head of the penis is grasped and the penis extended.

For transfusions, we used a 1-in., No. 20 needle, inserting it toward the symphysis pubis at about the midpenis position. A tough fibrous



# what makes *Donnatal*

so effective...  
so safe?...

Donnatal—one of the most widely employed spasmolytics today—derives its ever-increasing professional popularity from its dependable efficacy and its safety... demonstrated by controlled investigative studies,<sup>2,3,4,5,6</sup> and by gratifying clinical results in daily practice.

The secret of its success lies in its employment of only natural belladonna alkaloids,<sup>1,2,4</sup> in precisely and optimally balanced ratios—together with a small content of phenobarbital, for relief of the psychogenic factor by central and peripheral nervous sedation.

**A. H. ROBINS CO., INC. • RICHMOND 20, VA.**  
*Ethical Pharmaceuticals of Merit since 1878*

Hypertonicity, hypermotility and spasm of smooth muscle of the gastrointestinal, biliary and genito-urinary tracts. Donnatal is often effective in angina pectoris, hypertension, hyperacidity, parkinsonism and motion sickness.



TABLETS • CAPSULES • ELIXIR

...NATURAL

BELLADONNA ALKALOIDS

...BALANCED WITH

LABORATORY PRECISION



#### Tablets or Capsules:

1 to 2, three or more times daily (up to 9 tablets or capsules may be given within 24 hours without toxic effects).

**Elixir:** Infants: ½ teaspoonful 2 or 3 times daily, as necessary. Children: 1 teaspoonful, 3 or 5 times daily, as needed.

**Adults:** 1 or 2 teaspoonfuls, 3 or 4 times daily.

<sup>1</sup> Goodman, L. and Gilman, A.: *The Pharmacologic Basis of Therapeutics*, The Macmillan Co., New York, 1941. <sup>2</sup> Kilstein, S. J.: *Rev. Gastroenterol.*, 14:171, 1947. <sup>3</sup> Lee, L. W.: *Northwest State Med. J.*, 24:59, 1949. <sup>4</sup> Morrissey, J. H.: *J. Urol.*, 57:435, 1947. <sup>5</sup> Ricci, J. V.: *Contributions from Dept. of Gynecology, City Hospital, New York*, 1946. <sup>6</sup> Stephens, G. K.: *J. Oklahoma St. Med. Assoc.*, 42:246, 1949.

Rx



## CORRESPONDENCE

sheath surrounds each corpora so that some pressure is necessary before entering. When the corpora is entered, blood returns in the syringe or tubing as in any intravenous puncture. The needle and tube are strapped first to the end of the penis and then to the abdomen or leg.

In most cases no anesthesia was required, but we did abolish most of the pain when necessary with a small amount of 2% procaine just under the skin.

In infants a smaller gauge needle was used and the injection made with a syringe with a three-way stop-cock, just as is used in regular head vein sets.

M. D. BENTLEY, M.D.  
Sebewaing, Mich.

### Exercise Reprints Available

TO THE EDITORS: Would you kindly send me a reprint of "Postpartum Exercises" by Lt. Willie Rebecca Harvey, WMSC, which appeared in your August 15, 1949 issue.

H. B. WOOLLEY, M.D.  
Idaho Falls, Idaho

Because of the demand, a new supply of reprints of "Postpartum Exercises" has been ordered. A limited number are available again.—Ed.

### Reserves Copy of Index

TO THE EDITORS: I am one of the fortunate ones who regularly receive your journal. I am delighted to learn that a *Modern Medicine Index* of 1949 is being published. Would you please reserve a copy for me. Thank you for your wonderful service.

MORRIS FRANKLYN H. LEVY, M.D.  
Brooklyn

### Good Results with Blastomycosis

TO THE EDITORS: I have just read Dr. David T. Smith's nice article on immunologic types of blastomycosis (*Modern Medicine*, Jan. 1, 1950, p. 53).

During my absence several years ago, my locum tenens was treating with potassium iodide a patient who had a severe case of blastomycosis. The patient was fast getting worse. When I returned I suggested that we give him antimony and potassium tartrate intravenously. We did so, giving him  $\frac{3}{8}$  gr. for the first dose and  $\frac{3}{4}$  gr. thereafter three times weekly. Then, after 6 doses, we gave him  $1\frac{1}{2}$  gr. in the same routine. In about a month he was practically well, and in six weeks was completely recovered.

I have been impatiently waiting for another case. None have shown up in my practice. So far as I know, this treatment is original with me, and I am very anxious to have some of my fellow colleagues try it. The results, of course, may be just a "flash in the pan," so be patient with the suggestion.

Last month, I suggested antimony and potassium tartrate in the treatment of a suspicious inflammatory mass in the lung diagnosed as a possible malignancy or as an inflammatory tumor the size of a small orange. The patient also had a severe cough of a year's duration with profuse expectoration of muco-pus containing streptococci and staphylococci but no tumor cells. We fluoroscoped the lung three days ago. Cough, expectoration, and tumor are entirely gone.


F. A. ALLIN, M.D.  
San Antonio

*available* **NOW**  
**VITAMIN B<sub>12</sub>**

**FOR ORAL ADMINISTRATION**


**DOCIBIN**

Trademark



If your pharmacist does not  
stock **DOCIBIN**, he can order  
it from his wholesaler or from

**Walker** **VITAMIN PRODUCTS, INC.**  
MOUNT VERNON • NEW YORK



# C Chemoprophylaxis Against Pharyngeal Infections

## SAFE, TOPICAL METHOD

In Neiman's study\* on chemoprophylaxis with White's Sulfathiazole Gum conducted over a 9-month period on 199 medical students:

1. The incidence of primary pharyngitis in the treated group was *less than half* that in the controls. A less marked, but statistically significant, decrease was also observed in the incidence of colds and irritational pharyngitis.
2. "It is worthy of note that the mouths of over 100 persons were exposed to the drug in concentrated form daily for eight months, with *no untoward effects*."

As with the therapeutic use of Sulfathiazole Gum, the prophylactic application is safe because it is topical. In this series, for example, repeated examination of blood samples from unselected individuals in no case gave a positive test for sulfathiazole.

The dosage in these experiments was one to three tablets a day—an obviously economical procedure. No reactions were observed.

White's

## SULFATHIAZOLE GUM

### SAFE, TOPICAL CHEMOTHERAPY

Supplied in packages of 24 tablets— $3\frac{3}{4}$  grs. (0.25 Gm.) per tablet—sanitaped in slip-sleeve prescription boxes.

WHITE LABORATORIES, Inc., Pharmaceutical Manufacturers, Newark 7, N. J.

\*Neiman, I. S.: Prophylactic Value of Sulfathiazole, Archives of Otolaryn. 47:158-164 (Feb.) 1948.



**Patient under Treatment**  
**FOR URINARY**  
**TRACT INFECTION**



**P**rompt and effective relief from distressing symptoms of urinary tract infections often can be achieved through the action of orally administered Pyridium.

The analgesic action of Pyridium is entirely local, reducing the urinary frequency and pain and burning on urination, without systemic sedation or narcotic action.

Pyridium is virtually nontoxic in therapeutic dosage and can be administered concomitantly with streptomycin, penicillin, the sulfonamides, or other specific therapy.

Pyridium is the trade-mark of the Pyridium Corporation for its brand of Phenylazo-diamino-pyridine HCl. Merck & Co., Inc. sole distributor in the United States.



*The complete story of Pyridium and its clinical uses is available upon request.*

**Pyridium<sup>®</sup>**  
(Brand of Phenylazo-diamino-pyridine HCl)

**MERCK & CO., Inc.** Manufacturing Chemists **RAHWAY, N. J.**  
*In Canada: Merck & Co. Limited — Montreal, Que.*

## CORRESPONDENCE

### Long-sought Information

TO THE EDITORS: I will be ever so grateful to you if you could provide the address and apply it to the enclosed envelope for Dr. Charles G. Hutter, whose article on the bee cell pessary recently appeared in *Modern Medicine* (Jan. 1, 1950, p. 68).

Several years ago I inherited several of these pessaries from a preceptor. When my supply was gone I attempted to replenish it, but exhaustive inquiries produced no pessaries, only the information that the device had been manufactured in Germany and that none had been imported since before the last war and that no more would be available. You can imagine then what a very pleasant surprise it was to encounter the bee cell pessary in your magazine.

I have received *Modern Medicine* almost since its inception and have found a pleasant surprise in practically every issue. My thanks for your very modern *Modern Medicine*.

JOHN L. LANGOHR, M.D.  
Columbia City, Ind.

Dr. Hutter says that the pessary he described was ordered from the Bee Cell Company, Box 212, Buffalo 5, N.Y.—Ed.

### Quantity Just Right

TO THE EDITORS: I first contacted your journal when I came to this country for my internship. Since then, I have come to like and appreciate more and more your very excellent medical digest. I would like to congratulate you on the fine quality and just right quantity of all these reports.

ROBERT GEIGER, M.D.  
New York City

### Questions Value of Prenatal Care

TO THE EDITORS: Before the advent of prenatal care, most women bore children without needing any stitches for repair.

After elaborate prenatal care, modern obstetricians expect major tears as a rule and routinely do episiotomies which require several sutures for repair.

Calcium with calcium-stabilizing factors, administered during normal pregnancy, prevents the physiologic softening of the maternal pelvis and causes the premature calcification and hardening of the fetal bones in general and of the fetal skull in particular.

ALFRED ROSSKAMM ROSS, M.D.  
New York City

### Alcohol Capacity

TO THE EDITORS: I recently read the news item on alcohol capacity (*Modern Medicine*, Aug. 15, 1949, p. 88) which quotes Dr. Henry W. Newman as stating that:


The maximum amount of alcohol that can be consumed daily by a man of average weight is a quart of 100-proof liquor. This maximum may be achieved only by maintaining the blood alcohol concentration at a high level, since the rate of alcohol metabolism increases with blood alcohol concentration.

This may be true of the majority of persons. However, there are notable exceptions. I know of one person who was said to consume habitually 4 or 5 qt. of whiskey per day without apparent intoxication, and I personally knew a man here in Los Angeles back in 1934 who took little water but who drank 2 to 3 qt.

(Continued on page 24)

# RUTAMINAL\*

Ocular  
Fundus in  
Degenerative  
Vascular  
Disease  
Hypertension,  
Diabetes,  
Arteriosclerosis  
—note  
tortuous,  
blood vessels,  
areas of  
exudation,  
hemorrhagic  
areas.



Normal  
Ocular  
Fundus



\*In keeping with newer clinical findings, the rutin content of RUTAMINAL has been increased to 60 mg. per tablet (three times the former rutin content) at no increase in cost to the patient.

\*RUTAMINAL is the trademark of Schenley Laboratories, Inc. and designates exclusively its brand of tablets containing rutin, aminophylline, and phenobarbital.

the  
protection  
of  
**rutin**<sup>1</sup>  
the  
action  
of  
**aminophylline**  
the  
sedation  
of  
**phenobarbital**  
—for  
use  
in  
selected  
cardiovascular  
and  
diabetic  
conditions  
in  
which  
excessive  
capillary  
fragility  
presents  
a  
complicating  
hazard  
—bottles  
of  
100  
tablets

**schenley** laboratories, inc., 350 fifth ave., new york 1, n. y.

© Schenley laboratories, inc.



# RIASOL in




BEFORE USING RIASOL

## THOUSANDS PRESCRIBE RIASOL

RIASOL CONTAINS 0.45% MERCURY CHEMICALLY COMBINED WITH SOAPS, 0.5% PHENOL AND 0.75% CRESOL IN A WASHABLE, NON-STAINING, ODORLESS VEHICLE

APPLIED LOCALLY—SIMPLE TO USE

# PSORIASIS



**AFTER USING RIASOL**

**SEND FOR A CLINICAL PACKAGE**  
**PROVE RIASOL YOURSELF**  
**SHIELD LABORATORIES**

12860 MANSFIELD AVENUE DETROIT 27 MICHIGAN

## CORRESPONDENCE

of "Old Quaker" per day without showing evidence of intoxication.

This man was approximately forty-five years old, 5 ft. 6 in. tall, weighed about 175 lb. He kept several bottles of "Old Quaker" in his desk, and through the day he would frequently pour out and drink an ordinary water glass full of whiskey. I know that it was real whiskey because I saw the seals removed from the bottles and tasted the liquor. Five minutes or less after taking a drink one could not smell it on his breath!

Too many workers try to oversimplify the question of intoxication.

The amount of alcohol in the brain is what determines intoxication. Building up the alcohol content of the brain is not so simple. The alcohol must be carried to the brain by the blood after absorption from the gastrointestinal tract. It is a matter of common knowledge that an individual often varies in his susceptibility to alcohol from time to time, and particularly that alcohol taken on an empty stomach is usually more apt to intoxicate than that taken after, or with, a full meal.

There is good reason to believe that absorption of alcohol from the gastrointestinal tract in different individuals varies considerably. This being true, only that portion absorbed would appear in the blood.

After absorption, alcohol in the blood is [1] temporarily stored in the tissues, including the brain, [2] oxidized, and [3] eliminated unchanged through the kidneys and lungs. These factors also are probably subject to rather wide variations.

Obviously, during the absorptive stage, the blood alcohol will be high-

er than the brain alcohol and, during the eliminative stage, with no more alcohol being absorbed, the blood alcohol will be lower than the brain alcohol. This was well shown by a pathologist in Milwaukee, whose name I have forgotten, about ten or twelve years ago. This fact also lessens the value of blood alcohol determination in medicolegal cases. It also renders of even more doubtful value in medicolegal cases the amount of alcohol found in the urine or expired air of a person.

I do not believe that any well-informed person will question the fact that alcoholic intoxication is a direct function of the percentage of alcohol in the brain. However, if a person absorbs alcohol slowly from the gastrointestinal tract and rapidly excretes and oxidizes that which is absorbed, there will be little for the brain to absorb. Another may absorb rapidly, excrete and oxidize slowly, and thus quickly acquire a high concentration in the brain, with consequent quick, intense intoxication.

From many years of observation of man, his drinking habits and reactions, and from thousands of autopsies, I am convinced that there are infinite variations of the absorption, excretion, and oxidation pattern in man and that anyone who makes categorical statements on any one phase of the problem opens his work to serious challenge. I make only one exception to this statement, and that is the well-established, positive correlation between alcoholic intoxication and the percentage of alcohol in the brain.

JOHN H. SCHAEFER, M.D.

Los Angeles



## "valuable therapeutic agent in dermatologic practice"

### 4 excerpts from the literature on Vioform

Overton

"The effect of 3% Vioform in a water-miscible base or in petroleum jelly has been assessed under carefully controlled conditions on a variety of dermatoses in 176 patients. It proved a useful local application in the treatment of the following conditions: coccogenic sycosis barbae, seborrhoeic dermatitis, otitis externa, acute vesiculo-papular eczema. The incidence of cases of intolerance was low."<sup>1</sup>

Sulzberger & Baer

"Both long clinical experience and the present studies indicate that Vioform (5-chloro-7-iodo-8-hydroxyquinoline) is a valuable remedy in topical therapy. . . . Since submitting this report we have confirmed our observations in hundreds of additional cases. The results have strengthened our conviction that Vioform preparations, while not panaceas, are among the most valuable local therapeutic agents with very low irritancy and a low index or potential of sensitization."<sup>2</sup>

Martin-Scott

"... in at least 37 cases in which penicillin had failed Vioform produced the desired result. . . . It is concluded that 3% vioform is a valuable addition to the medicants at present used in this country for pyococcal dermatoses."<sup>3</sup>

Perlman

"Its results in the treatment of infantile eczema, as well as in the eczemas of older children, have been found good . . . in fact, on occasions Vioform has been proved to be the topical answer to eczema when orthodox treatment with the tars failed."<sup>4</sup>

1. Martin-Scott, I.: Brit. Med. J., May 14, 1949. 2. Overton, J.: Brit. Med. J., May 14, 1949. 3. Sulzberger & Baer: Arch. Derm. & Syph., 58: Aug. 1948  
4. Perlman, H. H.: J. of Pediat., July 1948

cream

3% in a special water-washable (flesh color vanishing cream) base—50 Gm. jars and 1 pound jars.

**Vioform**

ointment

3% in a petrolatum base (tends to stain, should be covered in use)—50 Gm. jars and 1 pound jars.

**Ciba**

PHARMACEUTICAL PRODUCTS, INC., SUMMIT, NEW JERSEY

VIOFORM (brand of iodochlorhydroxyquinoline)

Trade Mark Reg. U.S. Pat. Off. 2/1970M

Increases Load Capacity by **100%**

... accommodates 2 instrument trays instead of one

## The "American" All-Purpose JUNIOR AUTOCLAVE (MODEL 8816)

- Fully automatic and self-compensating for all types of loads.
- Requires no valuable floor space. May be placed on any flat surface and operated from convenient electrical outlet.
- This all-purpose unit with 8"x8"x16" chamber will sterilize instruments, dressings, rubber gloves, solutions, antibiotics and all allied loads.
- The unit is automatically "burn-out-proof" and operates by one control lever which governs the complete functional cycle.

Presents a radical departure in autoclave design that meets every requirement wherever a single, pressure steam sterilizer of small size is indicated. Duplicates in every respect the efficient, precision performance of standard size "American" Surgical Supply Sterilizers widely used in hospitals.



Diagrammatic comparison of Model 8816 and 8"x16" cylindrical type. Note accommodation of 2 instrument trays instead of one.


Brings Hospital Sterilizing  
Efficiency to the  
**PROFESSIONAL OFFICE**  
**LABORATORY - CLINIC**



**SPACIOUS, DOUBLE-CABINET MODEL DB-16** is ideally constructed to accommodate **MODEL 8816**, thus combining the essentials of a practical Surgical Supply for the private office.

WRITE TODAY for complete information

**AMERICAN STERILIZER COMPANY**  
Erie, Pennsylvania



# KOAGAMIN<sup>®</sup>

IS INDICATED IN

## EVERY CASE

**REGARDLESS OF THE UNDERLYING CONDITION**

Its prompt action—a matter of minutes—differs from that of vitamin K and its synthetics, which must first be converted to prothrombin in the liver—a matter of hours.

Clinical investigation has demonstrated that vitamin K is useful only in cases where prolonged prothrombin time is a factor. Yet, even in these conditions, KOAGAMIN should also be used for its rapid action.

**KOAGAMIN IS INDICATED IN EVERY CASE!**

### PREOPERATIVELY—

provides a clearer field of operation—minimizes cauterization or need for local hemostatics.

### POSTOPERATIVELY—

for control of secondary bleeding.

### THERAPEUTICALLY—

aids in the control of bleeding and should be used routinely in blood dyscrasias and hemorrhagic conditions.

Supplied in 10 cc. diaphragm-stoppered vials.

Literature upon request.



CHATHAM PHARMACEUTICALS, INC.  
NEWARK 2, NEW JERSEY, U. S. A.

Available Through Your Physician's Supply House or Pharmacist



# Forensic Medicine

COMPILED BY ARTHUR L. H. STREET, LL.B.

**PROBLEM:** In a personal injury suit, did the trial judge properly refuse to permit a physician to testify over plaintiff's objection what plaintiff's admission record at the hospital showed concerning the cause of a previous hospitalization in another state and under another doctor's care? The record had been prepared from information given by plaintiff.

**COURT'S ANSWER:** Yes.

The Colorado Supreme Court decided that under the statutes of that state, which resemble those in force in most other states, "a physician without the consent of his patient is prohibited from disclosing any information however acquired in attending him" (210 Pac. 2d 214).

**PROBLEM:** In a malpractice suit it appeared that Dr. P, a cancer specialist, advised biopsy of an enlarged axillary lymph node. While the patient remained anesthetized, the specimen was examined in the hospital laboratory and a diagnosis of carcinoma was made. The woman's right breast was immediately removed by Dr. P, with Dr. H, resident hospital surgeon, assisting. When the operation was almost completed, the laboratory changed the diagnosis to Hodgkin's disease or lymphoma. Was the resident doctor liable for damage to the patient?

**COURT'S ANSWER:** No.

The California District Court of Appeal decided that defendant was not liable for the "most lamentable error" that resulted in serious in-

jury, because there was no proof that he had committed any wrongful or negligent act. The defendant simply followed instructions of Dr. P, who determined the necessity for the operation and the time and extent of the performance. The mistake had originated in the pathologic laboratory and a jury had exonerated the pathologist in a former trial of the case. In the meantime, Dr. P had died. In short, the appellate court decided that Dr. H, as an assistant, could not be held liable for the laboratory's mistake or Dr. P's hasty decision (211 Pac. 2d 6).

¶It appears that although courts generally hold a directing surgeon liable for damage resulting from neglect of assistants, selected or adopted by him, the assistant is not liable for the principal surgeon's fault—A.L.H.S.


**PROBLEM:** A dispute arose in a personal injury suit as to whether the plaintiff's disability was partly due to prior disease and injuries. Could the trial judge require plaintiff to consent to defendant's inspection of medical records covering prior treatment of plaintiff for such disease and injuries, despite a local statute disqualifying physicians from testifying without the patient's consent about confidential information obtained in treating a patient?

**COURT'S ANSWER:** Yes.

The Wisconsin Supreme Court reasoned that plaintiff should be requir-

(Continued on page 32)





**a newly  
accepted therapy**

Mounting clinical evidence, now accepted by the council on Pharmacy and Chemistry of the American Medical Association, continues to support the claims made for the efficacy of Resinat. The most recent studies, for example, demonstrate that complete symptomatic relief occurs in from 48 to 72 hours and is accompanied by regression of the ulcer crater in from two to four weeks, as seen in most of the 120 patients treated with Resinat.<sup>1</sup>

Resinat acts as an adsorbent which effectively neutralizes excess gastric acidity. It does not cause constipation nor does it produce acid rebound or other objectionable side effects.

Resinat is available in capsules (0.25 Gm.), Tablets (0.5 Gm.), Powder (1 Gm.).

1. Weiss, S., Espinal, R. B. & Weiss, J.: Therapeutic Application of Anion Exchange Resins in the Treatment of Peptic Ulcer, Review of Gastroenterology, 16:501-509, June, 1949.

*Literature and samples available.*



**THE NATIONAL DRUG COMPANY**

*More than Half a Century of Service*




**PHILADELPHIA 44, PA.**

*to the Medical Profession.*



*Psoriasis — duration  
15 years*



*Photographed after 8 weeks  
treatment with Mazon*

*Mazon and Mazon Soap act quickly and  
bring skin infections\* under control*

\*These infections include


**Eczema**  
**Psoriasis**  
**Alopecia**  
**Dandruff**  
**Ringworm**  
**Ivy Poisoning**  
**Athlete's foot**

The fundamental principles employed in the development of Mazon—careful selection of ingredients, skillful blending for ease of application, absorbability and patient acceptance—promote the successful treatment of obstinate skin conditions.

The distinctive characteristics and therapeutic value of Mazon have been demonstrated clinically for over 25 years. Skin irritations not caused by or associated with systematic or metabolic disease or disorders are usually amenable to treatment with Mazon.

# MAZON

**BELMONT LABORATORIES,**  
Philadelphia, Pa.



## Excel-Print<sup>\*</sup> STATIONERY

The stationery that costs only a trifle more than the plain printed but which only experts can tell from the genuine engraved.

- ▼ SUPERIOR QUALITY
- ▼ CORRECT STYLES
- ▼ PROFESSIONAL DIGNITY

ILLUSTRATED: Letterheads, 5½" x 8½", and envelopes, 3½" x 6", to match. Both "Excel-Printed". Our regular price: \$10.90 for 1000 of each, delivered FREE anywhere in the U.S.A.

# Stationery of Distinction

WITH 22 YEARS OF REPUTATION

### Free Samples and Catalogues

Samples of "Excel-Print" and plain print stationery and copy of BIG catalogue illustrating, describing, and pricing ALL items used in doctors' offices are yours on request. No obligation.

\*T. M. Reg. U. S. Pat. Off.

"Excel-Print" stationery is made only by us. It is a superlative product that combines an exclusive process with fine papers to create stationery that is in keeping with the dignity of your profession. You must see and inspect it to appreciate its true quality and consequent popularity. And when you compare its prices to what you now pay, or with what others are charging, you positively will not believe them to be true. Twenty-two years of organizational know-how, the finest of modern equipment and a volume that services more than 70,000 doctors, insures these statements. Let us give you all the facts.

PROFESSIONAL

PRINTING CO., INC.

America's Largest Printers to the Professions

202 TILLARY ST., BROOKLYN 1, N. Y.

STATIONERY • MISCOUNT PRODUCTS  
PRINTING • RECORDS • FILES & SUPPLIES

PROFESSIONAL PRINTING CO., INC.  
202-208 Tillary St., Brooklyn 1, N. Y. 2-3-0

Please send me samples of stationery and copy of your BIG General Catalogue.

Dr. \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

# ONLY THOROUGHbred

**JUMPING-JACKS®**  
FLEXIBLE SHOES FOR HARD WEAR


carry this  
BRAND



## THE JUMPING-JACK

brand is a guarantee of finest quality leathers and painstaking workmanship. Jumping - Jack's patented principle assures helpful support to young ankles.

FROM CRADLE TO 4 YEARS



**VAISEY-BRISTOL SHOE CO., INC.**

ROCHESTER 3, NEW YORK

MADE IN CANADA BY THE SAVAGE SHOE CO.,  
LIMITED • PRESTON, ONTARIO

ed to furnish to defendant all pertinent information concerning his previous disability, whether that information be possessed by him or under the control of another. As to records of a Veterans' Hospital, he could not be regarded as having control over them, but he should be required to consent to their examination by defendant's attorney or physician. The court required that he consent to inspection of the hospital records, nurses' and technicians' notes and records, attending physicians' reports, and so forth, so far as such records pertained to the prior disability.

The opinion recognizes limitations of the statutory rule that medical information cannot be disclosed without patient's consent. It is noted that the reason for the rule is that if a physician could testify without the patient's consent, the patient would be apt to conceal "vicious or uncleanly habits necessary for a physician to know in order to treat them properly" rather than be humiliated by their disclosure. If the disclosure will not "subject the patient to shame or affect his reputation or social standing, there is . . . sound reason why in the interest of truth and justice he should be compelled to disclose them. The physician's exemption from disclosure should be limited to such disclosures as would injure the patient's feelings or reputation" (39 N. W. 2d 675).

The Wisconsin court referred to its earlier decision in another case in which it was decided that, despite the statute, the following types of evidence were admissible in a court trial without the patient's consent: [1] testimony of a nurse assisting an attending physician, [2] hospital record made by a nurse on the



*in Mixed  
Bacterial  
Genitourinary  
Infections*

Aureomycin is now rapidly becoming recognized as a drug of choice in the treatment of mixed bacterial genitourinary infections, particularly those in which *Escherichia coli* and *Aerobacter aerogenes* play a part. Intractability of a genitourinary infection is an especial indication for aureomycin.

Aureomycin has also been found highly effective for the control of the following infections: African tick-bite fever, acute amebiasis, bacterial and virus-like infections of the eye, bacteroides septicemia, bouton-neuse fever, acute brucellosis, Gram-positive infections (including those caused by streptococci, staphylococci, and pneumococci), Gram-negative infections (including those caused by the *coli-aerogenes* group), granuloma inguinale, *H. influenzae* infections, lymphogranuloma venereum, peritonitis, primary atypical pneumonia, psittacosis (parrot fever), Q fever, rickettsialpox, Rocky Mountain spotted fever, subacute bacterial endocarditis resistant to penicillin, tularemia and typhus.

## AUREOMYCIN HYDROCHLORIDE LEDERLE

**Capsules:** Bottles of 25, 50 mg. each capsule. Bottles of 16, 250 mg. each capsule.

**Ophthalmic:** Vials of 25 mg. with dropper; solution prepared by adding 5 cc of distilled water.

# AUREO- MYCIN

LEDERLE LABORATORIES DIVISION *AMERICAN Cyanamid COMPANY*  
30 Rockefeller Plaza, New York 20, N. Y.



*when mental depression and nutritional inadequacy  
manifest themselves as*

**apathy**

**lethargy**

**physical debility**

**R<sub>x</sub>** 'Dexedrine' plus essential B vitamins

## **Theptine**

*a light and palatable antidepressant  
and restorative elixir*

Each 5 cc. (1 teaspoonful) contains:

'Dexedrine'\* Sulfate, 2.5 mg.; thiamine hydrochloride, 5.0 mg.;  
riboflavin, 0.45 mg.; niacin, 6.7 mg.

**Smith, Kline & French Laboratories, Philadelphia**

\*T.M. Reg. U.S. Pat. Off. for dextro-amphetamine sulfate, S.K.F.

Easter Island Figurine; Photo courtesy University of Pennsylvania Museum



patient's admission, used by the attending physician, and [3] testimony of an x-ray technician, including a plate made by him under the doctor's direction (276 N. W. 300).

¶Although courts in other states might agree with the conclusions of the Wisconsin Supreme Court, the summary of the decisions cited above should not be relied upon as necessarily reflecting the law in other states. For example, many courts might not concur in the view that, in the absence of specific statutory declaration, the confidential character of information acquired by a doctor from a patient is limited to information the disclosure of which would *humiliate* the patient.—A.L.U.S.

**PROBLEM:** After a woman had been returned to her room following a total hysterectomy, insertion of a catheter brought a discharge of mixed blood and urine. The patient and her husband both testified that, immediately after the operation, the defendant surgeon admitted that he had "nipped" the patient's bladder; the surgeon denied having made such a statement. The patient, who complained of burning, urgency, and incontinence after the hysterectomy, was examined by a urologist who found infection and calculi of the bladder but no surgical damage to the walls. Three months later, the urologist reexamined the patient and found a vesicovaginal fistula, which was later repaired by another doctor. Without expert medical testimony could a jury conclude that the first doctor had failed to use due care and skill?

**COURT'S ANSWER: No.**

Said the Ohio Court of Appeals: "We find no medical evidence relating to the nipping of the bladder as a probable cause of the burning, urging and incontinence, . . . or that there was any causal connection between such claimed nipping and the vesicovaginal fistula which was subsequently found" (88 N. E. 2d 76).



# PERTUSSIN

in successful use  
for over 30 years for  
**COUGHS** in

- Acute and Chronic Bronchitis
- Paroxysms of Bronchial Asthma
  - Whooping Cough
  - Dry Catarrhal Coughs
  - Smoker's Cough

In Pertussin—the active ingredient—Extract of Thyme (unique Taeschner Process) effects relief of coughs not due to organic disease, because it:

1. Relieves dryness by stimulating tracheobronchial glands.
2. Facilitates removal of viscid mucus.
3. Improves ciliary action.
4. Exerts a sedative action on irritated mucous membranes.

Pertussin is entirely free from harmful drugs of any kind. It is well tolerated by adults and children and is pleasant to take. It has no undesirable side action.





**NOW!** *Completely Safe...*

**GUSTAMATE**

BRAND OF GLUTACINATE

**NEW SEASONING AGENT ADDS TASTE  
APPEAL TO LOW-SODIUM DIETS . .  
WINS PATIENT-COOPERATION**

**GUSTAMATE\*** is a unique, new, nonmineral seasoning agent, completely safe for routine use in low-sodium diets. Its principal component is monoammonium glutamate, with balanced proportions of the amino acids, glycine and glutamic acid, established as harmless even when taken in quantities far in excess of the amounts provided in the average daily intake of **GUSTAMATE**.

Monoammonium glutamate is similar in flavoring effect to monosodium glutamate, long used in hotel and restaurant cuisines to bring out the natural flavors of foods. **GUSTAMATE**, however, contains no sodium.

**INDICATIONS:** In sodium-restricted diets prescribed for congestive heart failure, hypertension, renal disease, obesity, certain disorders of pregnancy (e.g., toxemia), and in conditions characterized by poor or finicky appetite.

**SUPPLIED:** As white, crystalline granules in salt-shaker-type dispensers containing 1 ounce. Available at leading pharmacies.

*Arlington*

**THE ARLINGTON CHEMICAL**

\*The word GUSTAMATE is a trademark of The Arlington Chemical Company.

# MINERAL-FREE SALT SUBSTITUTE..

## FEATURES OF GUSTAMATE

### *Ensures Safety*

- Free from sodium
- No other metallic ions
- No disturbance of mineral balance
- Contains substances normally participating in metabolic processes
- Can be used safely over long periods

### *Increases Palatability*

- Brings out the natural flavor of foods
- Enhances effect of other seasonings
- Often suppresses undesirable taste features
- Prolongs agreeable taste sensations
- Stimulates appetite and salivation

Complete literature on request

COMPANY . . YONKERS 1, . . NEW YORK

# Questions & Answers

*All questions received will be answered by letter directed to the petitioner; questions chosen for publication will appear with the physician's name deleted. Address all inquiries to the Editorial Department, MODERN MEDICINE, 84 South Tenth Street, Minneapolis 3, Minnesota.*

**QUESTION:** Recently a typical pituitary dwarf has come under my care. He is twenty-five years old. Roentgenograms reveal open epiphyses. Have you any suggestions for treatment?

M.D., California

**ANSWER:** *By Consultant in Internal medicine.* Theoretically, growth is possible if epiphyses are not united but, actually, results are disappointing, especially at the age your patient has attained. Extract of anterior pituitary gland, possibly combined with thyroid extract, is a logical therapy.

**QUESTION:** I have treated a patient with pulmonary tuberculosis with streptomycin for two years and have had good results. May treatment be continued without fear of bad reactions?

M.D., New York

**ANSWER:** *By Consultant in Chest Diseases.* It is impossible to give a specific answer to a question on tuberculosis in which the particulars of the case are not given.

Streptomycin is a valuable drug in the treatment of tuberculosis but has very definite limitations and must, therefore, remain part of a well-integrated program in which prolonged bed rest, the various collapse measures and even resection are carried out when indicated. After streptomycin therapy of three to four months,

most patients will have organisms that are resistant to streptomycin. It has recently been shown that para-aminosalicylic acid (PAS) given in combination with streptomycin delays emergence of resistant strains. Studies should be done to determine the sensitivity of the organisms.

Reactions to streptomycin therapy must still be carefully watched for, but they are relatively few when the drug is given in optimum dosage. Dosage, in a very general way, is 1 gm. of regular streptomycin, or up to 2 gm. daily when dihydrostreptomycin is being given. Recently new programs of interrupted therapy, injections every other day or only once weekly, have been attempted and show promise of giving good results with even less toxicity.

**QUESTION:** A healthy, twenty-eight-year-old woman noticed, about four months ago, that her breasts were becoming smaller. Her child was four years old. There have been no other physical changes. Laboratory tests, including study of ketosteroids, have been negative. Can you suggest a reason for the atrophy and possible treatment?


M.D., Ohio

**ANSWER:** *By Consultant in Gynecology.* Atrophy of the breast may result from hormonal changes or from loss of tissue response to hor-

*compatible with the*

# *deficiency theory*

*in the management of gastrointestinal disorders . .* **Viodenum**



"The treatment of . . . gastric ulcer and duodenal ulcer . . . is based on the supposition that the normal functions of the stomach and the duodenum are maintained by various biologically active substances, the absence of which favours ulcer formation."<sup>\*</sup>

<sup>\*</sup>Humbler, O., *Lancet*, **251**, 272 (1946).

"... not only were these patients relieved of their symptoms, but in all cases included in this particular report there was roentgenologic evidence of ulcers having healed . . . it is not expected that . . . duodenal . . . will prove to be a specific for peptic ulcer, for I do not believe that any single substance will ever be able to correct all the interacting factors responsible . . . I am fully convinced, however . . . the protecting mechanism inherent in duodenal . . . would be invaluable in the treatment of ulcer."<sup>\*</sup>

<sup>\*</sup>Rivers, A. D., *Am. J. Dig. Dis.*, **2**, 189 (1935).

"A consideration of the natural course of ulcerative colitis led to the theory that in some cases the condition might arise as the result of a deficiency. Preliminary investigations suggested that the missing hypothetical factor might be present in or produced by the intestine. Feeding experiments . . . showed that remissions could be induced regularly by giving uncooked pig's small intestine by mouth . . . the results obtained with this treatment do not appear to be coincidental or psychological; they are compatible with the deficiency theory advanced . . ."

<sup>\*</sup>Gill, A. M., *Lancet*, **2**, (1945).

"Duodenal substance [Viodenum] was administered to thirty-five patients who had chronic ulcerative colitis . . . no other specific medication was used . . . the results obtained in 85 per cent of the patients were very favorable . . . the majority of the patients gained weight . . . felt better and ate better . . . duodenal substance [Viodenum] may be considered a very valuable aid in the therapy of chronic ulcerative colitis."<sup>\*</sup>

<sup>\*</sup>Streicher, M. H., *J. Lab. Clin. Med.*, **33**, 1633 (1948).

RAW duodenum desiccated and defatted at body temperature  
Provided in powder or ten grain tablets:

Literature available upon request

**Viodenum**

**VIOLIN LABORATORIES**  
MONTICELLO, ILLINOIS • U. S. A.



# BIRTCHE

## HYFRECATOR

Used by More Than

### 60,000 DOCTORS

... for the removal of skin growths, tonsil tags, cysts, small tumors, superfluous hair, and for other technics by electrodesiccation, fulguration, bi-active coagulation.

Now, completely redesigned the new HYFRECATOR provides more power and smoother control... affording better cosmetic results and greater patient satisfaction. Doctors who have used this new unit say it provides for numerous new technics and is easier, quicker to use.

Send for descriptive brochure, "Symposium on Electrodesiccation and Bi-Active Coagulation" which explains the HYFRECATOR and how it works.



Send for Free Literature

To: The BIRTCHE Corp., Dept. A 3-0  
3087 Huntington Dr., Los Angeles 32, Calif.

Please send me free booklet, "Symposium on Electrodesiccation & Bi-Active Coagulation."

Name \_\_\_\_\_

Street \_\_\_\_\_

City \_\_\_\_\_

State \_\_\_\_\_

monal stimulation. The case under discussion is apparently an example of postlactation atrophy. If the patient is still menstruating normally, the atrophy is probably due to change in the breast, not in the hormone system. If such is true, hormone therapy would not be of benefit. If, however, there is a coexisting amenorrhea, estrogen therapy given in cyclic manner may stimulate growth of the breasts.

**QUESTION:** A man of forty, in good health, has consulted me for almost constant perspiration on the back. Palms or other parts of the body are not affected. A thorough examination reveals nothing significant. What is your opinion of the cause of the disturbance?

M.D., New York

**ANSWER:** By Consultant in Dermatology. Since the patient has had a thorough examination, it can be assumed that systemic disease is not a factor in this case. Localized disturbances of perspiration are usually related to structural or functional disturbance of nerve supply to the area. Such is probably the case in this instance.

**QUESTION:** Has a new drug been advanced for trigeminal neuralgia?

M.D., New York

**ANSWER:** By Consultant in Neurology. No specific new treatment for trigeminal neuralgia has been devised. The drug now being used is trichlorethylene, a liquid which the patient drops on a piece of cotton and inhales. A large number of patients with trigeminal pain are relieved by this drug. Occasionally, large doses of vitamin B<sub>1</sub> and C have proved helpful. The best treatment, when pain is severe, is section of the fifth cranial nerve.

**casimir funk laboratories, inc.**  
**announces**

# **pregnenolone**

**ENELONE** (trade mark)

**for experimental  
clinical investigation in  
arthritis**

Pregnenolone is a steroid substance which in limited experimental work has been demonstrated to have therapeutic activity approaching that of cortisone and ACTH in the treatment of rheumatoid arthritis . . . effective both *orally* and *parenterally* — and with early promise of plentiful supply.

Casimir Funk Laboratories is now making pregnenolone available to rheumatologists, clinicians, hospitals and others competent and eligible in conducting such clinical experimentation and investigation under the Federal Food, Drug and Cosmetic Act.

Available sizes:

**ENELONE oral**  
tablets of  
pregnenolone acetate

50 mg. tablets  
100 mg. tablets

**ENELONE injectable**  
aqueous crystalline suspension  
of pregnenolone

100 mg. per cc., 9 cc. vial

Inquiries should be addressed to . . .

Director of Research

**casimir funk laboratories, inc.**

affiliate of U. S. Vitamin Corporation

250 east 43rd street • new york 17, n. y.



When you prescribe iodine

# *Specify* **ORGANIDIN<sup>®</sup>**

**A Demonstrably Superior  
Iodine Preparation**

- WELL TOLERATED
- HIGHLY EFFECTIVE

*Organidin* (Wampole) is an exceptionally well tolerated, stable preparation of iodine organically combined by reaction with glycerin for internal administration.

Clinical studies<sup>1</sup> have demonstrated the superiority of *Organidin* to Lugol's Solution (inorganic iodine) at corresponding iodine levels for dosage periods of 6 to 30 days. *Organidin* may be administered at high dosage levels and for prolonged periods.

With *Organidin* iodine absorption is prompt, side effects are infrequent and mild. Whenever systemic iodine therapy is indicated (for hyperthyroidism, prophylaxis of goiter, syphilis, and cardiovascular conditions, or as an expectorant for relief of asthma or bronchitis) prescribe stable, well tolerated *Organidin*. Supplied in bottles of 30 cc. with dropper designed to deliver approximately 1 minim (7.5 mg. iodine) per drop. *Samples and literature on request.*


1. *Blaugher, D.: South Dakota J. Med. & Pharm., 1:425, 1948*

**HENRY K. WAMPOLE & CO., INCORPORATED**

*Manufacturing Pharmacists Since 1872*

PHILADELPHIA 23, PA.





**Chronic osteomyelitis** of 12 years. 14 surgical procedures had failed to close cavity. Pain and foul discharge caused patient to request amputation.



**Treatment with Chloresium** brought progressive closure. Drainage and odor stopped. Pinch grafts at base were successful and cavity closed completely.

## When tissue healing is the problem use Chloresium Therapy

**Clinically proved . . . More than 1150 cases reported in medical journals**

Clinical results in over 1150 cases with Chloresium, the therapeutic water-soluble chlorophyll preparations, have now been reported in the literature by eminent clinicians.\* Complete healing has been the rule in the great majority of these cases, although most of them had previously failed to respond to other methods of treatment.

The remarkable results obtained with Chloresium preparations are

# Chloresium

**Therapeutic chlorophyll preparations**  
**Solution (Plain); Ointment; Nasal**  
**and Aerosol Solutions**

**Ethically promoted—at leading drugstores**

U.S. Pat. 2,120,667. Other Pats. Pend.

due to the therapeutic action of the water-soluble derivatives of chlorophyll. They are natural biogenic agents which accelerate normal cell regeneration, thus measurably hastening the healing process. At the same time, they help control superficial infection, provide symptomatic relief and deodorize foul-smelling suppurative conditions.

\*Complete bibliography and reprints available on request.

### FREE—CLINICAL SAMPLES


**RYSTAN CO., INC., Dept. MM-2**  
7 N. MacQuesten Pkwy., Mt. Vernon, N. Y.

Please send clinical samples of Chloresium Ointment and Solution (Plain). I also want literature ☐; samples of Nasal and Aerosol Solutions ☐ (check if desired).

Dr. \_\_\_\_\_

Address \_\_\_\_\_

City \_\_\_\_\_ Zone \_\_\_\_\_ State \_\_\_\_\_



# Delicious Hard Candy and

•

• •

20,000  
units  
of  
penicillin

## PONDETS\* PENICILLIN TROCHES

**For local therapy and prophylaxis of oral infections caused by penicillin-sensitive organisms.**

Delightful tasting—welcomed by young or old. Potent—supplies 20,000 units penicillin in slowly dissolving hard candy base. Effectiveness lasts approximately one half-hour.

Trade Mark

*Wyeth* Incorporated, Philadelphia 3, Pa.



## *Special Article*

### Anesthetic Disasters

R. A. GORDON, M.D.\*

*Prepared for Modern Medicine*

A DANGEROUS and unjustified sense of security in the administration of anesthetics has arisen during recent years.

Despite the great advances in technic and management of anesthesia and the resulting decrease in the fatalities attributable to that agent, anesthetic disasters continue to take a considerable toll of life and health. These catastrophes are all the more tragic because nearly all could be avoided.

#### ASPHYXIA DURING ANESTHESIA

Most patients who die during anesthesia succumb to asphyxia. Death from this cause is in nearly every instance due to lack of technical ability and judgment on the part of the anesthetist.

Asphyxia may be due to actual obstruction of the respiratory passages, to depression of respiratory function to the point where the patient is no longer able to provide himself with adequate oxygen for survival, or to reduction of the oxygen content of the inhaled atmosphere below what is necessary for life.

Not all anesthetic calamities result in immediate death. Prolonged or severe anoxia frequently produces degenerative changes in the cerebral cortex which may terminate in death some hours or days after operation or cause mental changes

\* Clinical Teacher in Anesthesia, University of Toronto; Senior Anesthetist, Toronto General Hospital.

## *SPECIAL ARTICLE*

in the survivor. These sequelae may be avoided by maintenance of an adequate airway.

Under conditions of anesthesia, most patients require at least 20% of oxygen in the inspired atmospheric air. A lesser concentration than this, if imposed for longer than a very few minutes, may result in death or crippling mental changes. Some patients require more oxygen, particularly those who are anemic or in a state of shock or who have an increased metabolic rate because of fever or hyperactivity of the thyroid. Sound judgment on the part of the anesthetist is essential to prevent disaster in such cases.

Depression of the patient's respiratory function to the point of absolute or relative asphyxia is a matter of overdosage of the anesthetic agent. Overdosage in this respect must be considered in relation to the physical status of the patient and the preoperative use of sedative drugs. This is again a matter where good judgment is of paramount importance.

Some physicians do not realize that asphyxia from respiratory depression is possible while the patient still breathes. The act of breathing does not of itself insure adequate oxygenation. If the volume of respiratory exchange is not adequate to provide sufficient oxygen from the particular atmosphere which the patient is breathing, asphyxia results. Disaster from this cause may be avoided by artificial respiration and is inexcusable today.

### **OBSTRUCTION OF AIRWAY**

Asphyxia from blockage of the airway may be caused by simple obstruction by the tongue and epiglottis, the presence of foreign material in the airway, or laryngeal spasm. Other rare causes exist.

Many disasters have occurred because of a mistaken idea that movement of the chest is indicative of breathing. Actually, the individual who is being asphyxiated by occlusion of the airway makes violent respiratory efforts until the respiratory center finally fails because of anoxia. The only proof that breathing is really taking place is demonstration of the movement of air or anesthetic mixture into and out of the respiratory tract.

When a rebreathing reservoir bag or bellows is in use, respiration may be confirmed by movements of the bag or by

hearing or feeling the passage of air. Occlusion of the airway is a frequent cause of disaster during tonsillectomy when an endotracheal tube is not used.

Obstruction of the airway by foreign material is most frequently due to aspiration of vomitus or blood. The chances of aspiration of vomitus may be reduced by careful management of the induction period and by avoiding general anesthesia for patients who have recently taken food.

When this latter precaution is not possible, vomiting should be anticipated by the provision of adequate suction equipment and the induction of anesthesia with the patient on the side in the head-down position. A tight-fitting endotracheal tube should be used to prevent the aspiration of vomitus.

When vomiting has not been anticipated, adequate postural drainage and suction must be used to remove all vomited material from the airway.

Obstruction of the airway by inhalation of blood occurs only when surgery is performed within the airway. Most fatalities from aspirated blood are associated with tonsillectomies and adenoidectomies and may be absolutely avoided by the use of an endotracheal tube and a small gauze pack in the lower pharynx, coupled with proper positioning of the patient in the postoperative period so that blood drains from the mouth. When the anesthetist is unable to perform endotracheal intubation, great care must be taken to suck all blood from the pharynx promptly.

#### LARYNGEAL SPASM AND CONVULSION

Asphyxia due to laryngeal spasm occurs not infrequently. Disaster may be avoided by forcing oxygen into the respiratory tract as the glottis begins to relax. This relaxation always occurs before death from asphyxia supervenes, and the alert anesthetist is usually able to revive the patient by providing adequate oxygenation at the earliest possible moment. However, patients who are already in poor physical condition may not tolerate the period of anoxia that necessarily occurs with the spasm.

Laryngeal spasm may usually be prevented by a nonirritating induction of anesthesia, with the avoidance of stimulation of the larynx by secretions until full surgical anesthesia is

## SPECIAL ARTICLE

obtained, and by establishing and maintaining adequate surgical anesthesia before surgical stimulus is applied.

Convulsions during anesthesia are fortunately rare, but death following such an episode is also the result of asphyxia. The cause of such convulsions is yet obscure, but dehydration, fever, anoxia, and carbon-dioxide retention probably all contribute.

Control of the convulsion by an intravenous barbiturate and by adequate artificial ventilation with oxygen should prevent a major disaster.

### SPINAL ANESTHESIA

Death during spinal anesthesia is rarely due to an idiosyncrasy of the drug used. The usual cause is anoxia from deficient circulation coupled with diminished respiration because of paralysis of intercostal muscles.

This disaster may be avoided by proper control of the blood pressure through the use of vasopressor drugs and of parenteral fluid therapy and the administration of oxygen.

Among the disasters associated with spinal anesthesia, too little mention is made of meningitis resulting from infection and of permanent neurologic damage caused by mechanical and chemical factors. Scrupulous attention to the details of asepsis in the technic of inducing spinal anesthesia is essential.

Mechanical injury to the spinal cord rarely if ever occurs when the site of puncture is below the second lumbar interspace. Damage to nerve roots may be avoided if the lumbar puncture needle is kept in the midline.

Most instances of chemical damage to the central nervous system following spinal anesthesia may be traced to contamination of the drug with alcohol or some similar solution used to sterilize the ampule. Cracked and defective ampules cannot be entirely eliminated, but contamination is readily detected before the injection is made if ampules are sterilized in highly colored solutions.

### EXPLOSIONS

The most spectacular but least frequent accidents associated with inhalation anesthesia are explosions. All the agents commonly used for inhalation anesthesia will explode under

clinical conditions, with the exception of nitrous oxide, chloroform, and trichlorethylene.

Explosion of an anesthetic mixture may cause immediate death from disruption of the pulmonary alveoli or later death from burns in the respiratory tract. Such catastrophies may be prevented by precautions which prohibit the accumulation of static charges in operating rooms and the avoidance of explosive anesthetic mixtures when open flames and open electrical equipment are used.

## Aureomycin Therapy of Herpes Zoster

MAXWELL FINLAND, M.D., EDMUND F. FINNERTY, JR., M.D.,  
HARVEY S. COLLINS, M.D., JOHN W. BAIRD, M.D., THOMAS M.  
GOCKE, M.D., AND EDWARD H. KASS, M.D.\*

AUREOMYCIN is apparently specific for herpes zoster but should be given in the early or active phase.

If treatment is started within two weeks after the first eruption, Maxwell Finland, M.D., and associates of Harvard University and Tufts College, Boston, find that postherpetic pain can usually be prevented.

The drug was given for initial, acute, or late involvement in 24 cases. Cervical, thoracic, and lumbar segments were affected as well as branches of the trigeminal nerve. In a few instances lesions were generalized.

A total daily dose of 4 gm. is taken orally, 1 gm. after each meal and at bedtime, for two to four days or until lesions are well dried and encrusted. During the next three to five days, 0.5 gm. is administered four times daily.

The smaller dose may be given throughout the course when the regular amount is not tolerated. If preferred, 500 mg. is injected by slow intravenous drip in 500 or 1,000 cc. of 5% dextrose solution without alkaline buffer. The procedure usually takes about an hour.

Vesicles heal rapidly and completely. New lesions occasionally appear during the first day of treatment but not later unless the course is interrupted. Pain also subsides after the first day.

Aureomycin was not beneficial in 4 cases of herpes labialis. The lesions extended during therapy.

\* Aureomycin treatment of herpes zoster. *New England J. Med.* 241:1037-1047, 1949.



## Venous Stasis in the Legs

W. J. MERLE SCOTT, M.D., AND MICHAEL RADAKOVICH, M.D.\*

*University of Rochester, N. Y.*

**C**HRONIC ulcerative lymphedema of the legs may be diagnosed by a simple volumetric technic even when the superficial veins can be neither seen nor felt.

The lymphedema and venous stasis are effectively controlled by a pneumatic bandage which exerts uniform, rhythmic pressure by means of an inflatable bladder in a nonelastic sheath.


### VENOUS STASIS TEST

When varicose veins are concealed by induration of skin and subcutaneous tissues, ordinary tests of vascular function may be impossible. However, incompetence of perforating vessels is often shown by the differences in measurements of the leg

after changes in position, explain W. J. Merle Scott, M.D., and Michael Radakovich, M.D.

Enlargement is quickly and accurately estimated with a 25-foot cloth tape measure wrapped from ankle to knee.

The leg is measured first with the patient supine. After the limb has been raised at a 45° angle and held in that position for five minutes, the end of the tape is fastened with adhesive 2 in. above the external malleolus, and twenty parallel turns are made (Fig. a). The tape should rest against the skin without tension and the turns should not overlap. The terminal point directly in line with the initial point is marked and a direct reading is made from the tape.



\* Venous and lymphatic stasis in the lower extremities. *Surgery* 26:970-986, 1949.

The tape is removed and a tourniquet applied to occlude superficial veins (Fig. b). The subject stands for a second measurement (Fig. c). Tourniquet and tape are removed, and after five minutes of standing, the multiple circumferential measurement is again made (Fig. d). The first measurement is then repeated.

Venous stasis is indicated by an increase of more than 6 in., particularly on standing with the tourniquet. The greatest increase is always on the second measurement and may be as much as 20 in. If varicose veins are prominent and perforating veins competent, volume does not increase much until the tourniquet is removed.

#### LEGGING

Stasis is controlled by application of uniform pressure obtained with the Aero-pulse legging which has an outer casing of Cape Cod duck closed with a zipper. A butterfly-shaped rubber bladder fits into an inner pocket and is inflated through a tube and valve at the legging top. The hand pump used for inflation has a regulator which controls air pressure.

In most cases a level of about 35 mm. of mercury is adequate. The

balloon pulsates and pressure is raised by contraction of calf muscles during walking and reduced below the initial level by intervals of relaxation. Pulsation is applied uniformly over the entire lower leg. Exercise is beneficial rather than detrimental and ulcers may heal more rapidly than with bed rest.

If ulcers are below the malleoli, more pressure is sometimes needed in that area. A second small bladder can be inflated to a level of 50 mm. while the large container remains at 35 mm.

Bacterial or fungous dermatitis occasionally requires special treatment while the legging is worn or, in extreme cases, before the device can be applied. Perspiration is absorbed with a cotton understocking and, if necessary, by a layer of Cellu-cotton or an old Ace bandage loosely wrapped. An insulating layer of inert plastic or cellophane may be used if the patient is allergic to the rubber.

The support is not used when the leg is elevated. After circulation improves, the bandage can safely be removed during active periods of one to three hours daily. However, stasis develops and ulcers quickly reappear if treatment is discontinued.

**G**UAIAIC TEST FOR OCCULT BLOOD in the feces is suitable for office use and easy to employ. A preliminary meat-free diet is not required. In the technic used by Stanley O. Hoerr, M.D., William R. Bliss, M.D., and James Kauffman, M.D., of Ohio State University, Columbus, a bit of feces is smeared directly onto simple filter paper. Then 1 or 2 drops of each of the three reagents, guaiac solution, glacial acetic acid, and hydrogen peroxide, are added in sequence. Color change to blue or dark green within thirty seconds denotes presence of blood.

*J.A.M.A. 141:1213-1217, 1949.*

## Chilblains

JOHN T. INGRAM, M.D.\*

*General Infirmary, Leeds, England*

**N**o therapy is specific for chilblains, but empirical treatment, varied to suit the individual, will relieve all but a few patients.

Chilblains are an expression of an individual susceptibility to react to cold in a particular pattern. Impaired peripheral circulation predisposes to chilblains. The tendency is influenced by many different factors, internal and external, and may be affected by prophylactic measures, believes John T. Ingram, M.D.

### PROPHYLAXIS

Warm clothing, particularly on the extremities, prevents or modifies the effects of cold and humidity which precipitate vascular changes. Two pair of socks for men and elastic understockings for women may be advisable in winter. Clothing should not be tight and binding.

Exercise is important in the maintenance of circulatory tone. Feet or hands which have become cold should not immediately be exposed to warmth from fire or hot bottles.

### TREATMENT

Stimulation of circulation by electrical and radiation therapy, massage, and passive and active movements is often effective for chilblains.

An attack may be relieved by ultraviolet light therapy or by fractional

doses of x-rays, but these measures should not be employed by the inexperienced.

The only benefit of local stimulants appears to be derived from the accompanying massage. Irritation from local applications may cause dermatitis or blistering.

Calcium, calciferol, thyroid, and nicotinic acid may be helpful but the exact action of some of these drugs is not known. No scientific evidence supports the claims made for calcium, but patients who demand such treatment may obtain psychological benefit.

A high level of calciferol in the blood is not protective, but the metabolic shock of a massive dose may have some effect. If employed at all, this measure should be used with care and only for a short period.

Small doses of thyroid extract are often helpful by improving the circulation. Nicotinic acid acts as a peripheral dilator. Purgatives may abort an attack.

For cases which persist in spite of medical treatment, sympathectomy may be indicated. Otherwise, removal to a different climate may be the only solution.

### DIAGNOSIS

Most commonly, chilblains appear as an itching, erythematous, and congested superficial swelling on the fin-

\* Refresher course for general practitioners: chilblains. *Brit. M. J.* 4639:1284-1286, 1949.

gers and toes. The aspect may differ somewhat on the breast, buttocks, upper arms and lower legs, where the lesions are large and more diffuse. When the ear is involved, little swelling is possible, and burning, irritation, blistering, and atrophy or calcification occur instead.

The young, particularly at puberty, are most susceptible. Chilblains are rarely seen during pregnancy.

To be differentiated are:

► *Erythema nodosum*—This condition often follows a respiratory infection and is usually distinguished by tenderness, pain, and pyrexia.

► *Erythema induratum*—Lesions are more deep seated than those of chilblains. The skin may ulcerate, but does not itch.

► *Drug rash*—Eruptions are usually larger and more scattered than with chilblains and are symptomless.

► *Erythema multiforme*—Lesions are spread more profusely in a particular pattern and cause less irritation than with chilblains.

► *Lupus erythematosus*—Eruption is usually more defined than chilblains and appears on the backs of the fingers, commonly between the joints and about the nail folds.

**D**IAGNOSIS OF MUMPS is assured when gelatinous edema is found peripheral to enlargement of the parotid or submaxillary glands. If the margin of the edematous region is gently tweaked, the area quivers like jelly. A similar type of edema, not caused by mumps, is occasionally found in other soft tissues, as with insect bites or poison ivy. However, George Heller, M.D., of Englewood, N.J., has never seen a like swelling in any infection which might be confused with mumps.

*Am. J. Dis. Child.* 78:903-905, 1949.

**W**HOOPING COUGH is effectively treated by chloramphenicol. The antibiotic is usually administered orally, but Eugene H. Payne, M.D., Detroit, and associates of the Inter-American Coporate Service of Public Health, Cochabamba, Bolivia, find that results are equally good when the drug is given rectally as a suppository or intravenously as a solution in propylene glycol. Fever usually disappears during the second day of treatment and paroxysms on the third to sixth day. A light cough may persist for several days longer, probably because of residual inflammation in the tissues. One week after therapy, culture of cough plates of 50 children who had had severe involvement indicated that all were free of infection. The only side effect, noted in a few children under six months of age treated orally, was slight nausea, which may have been caused by the disagreeable taste. The total rectal dosage ranged from 1.5 to 4.5 gm., depending on the weight of the child.

*J.A.M.A.* 141:1298-1299, 1949.

## Brain Tumors in Children

A. EARL WALKER, M.D., AND THERON L. HOPPLE, M.D.\*

*Johns Hopkins Hospital, Baltimore*

*Toledo*

**I**NTRACRANIAL TUMORS found in childhood differ in symptoms, pathology, and course from those of adult life.

The majority arise along the medulla, pons, and third ventricle and soon block the exit of ventricular fluid, causing internal hydrocephalus. In a series of 100 cases with onset before the sixteenth birthday, most of the growths observed by A. Earl Walker, M.D., and Theron L. Hopple, M.D., lay beneath the tentorium.

Many lesions occurred during the sixth year, particularly medulloblastomas. No acoustic neuroma or pituitary adenoma was noted and meningioma was relatively rare.

Since a child's brain has great compensatory power, a tumor frequently produces no symptoms until large enough to cause intracranial hypertension by mass, blockade of ventricular fluid, or both. In babies and young children cranial sutures separate and the head enlarges.

At older ages the skull is rigid and the brain is molded into all crevices, replacing cerebrospinal and ventricular fluid, squeezing liquid from interstitial spaces, and constricting



the arteries. Cerebellar tonsils may herniate about the medulla oblongata into the occipital foramen and seriously depress vital functions.

The general prognosis for a child with brain tumor is rather poor, yet cerebellar astrocytomas and some of the cerebral tumors may be removed successfully.

As symptoms are often nebulous and misleading, all available diagnostic technics should be employed.

*Vomiting* occurs with or without nausea, sometimes suggesting gastrointestinal disease. The symptom is often precipitated by illness or a head injury and may stop suddenly for weeks, then start again with or without provocation.

*Headache* is not specific but is frequently referred to the frontal region, and occipital or suboccipital discomfort is common. Pain typically starts early in the morning, often with vomiting, which may give relief. Aggravation by stooping, coughing, or straining while at stool is rarely described.

*Staggering gait*, the third most frequent symptom, is likely to begin

\* Brain tumors in children. J. Pediat. 35:671-687, 1949.

gradually and be attributed to weakness from vomiting.

*Convulsions*, which often result from cerebral growths, resemble idiopathic epilepsy and are usually generalized, but may start with jerking of the face, arm, or leg. Posterior fossa tumors occasionally cause attacks of rigid opisthotonos, sometimes with respiratory and cardiac involvement.

*Sutural diastasis* of infants and small children may be palpated or recognized by a cracked-pot percussion note.

*Weakness* of an extremity, side of the face, or eyelids localizes the neoplasm, as a rule, but diplopia indicates generalized pressure affecting the abducens nerve.

In all suspected cases of brain tumor, roentgenograms of the skull should be made in anteroposterior and lateral stereoscopic views. The lesion may be located by open sutures, calcification, calvarial distortion, or erosion. If the site is in doubt, special films of the optic foramen are made. The wrist should be examined for a lead line.

Electroencephalography is done, chiefly to show cerebral tumors and exclude epilepsy.

In older children, both site and type of growth may be determined by angiography; the common carotid artery, occasionally the vertebral artery, can be punctured through the skin. Vessels of babies may be inaccessible.

Pneumoencephalography is useful and safe if intracranial pressure is not elevated. With high pressure, ventriculography is preferable but may cause circulatory changes unless the tumor is immediately removed.

Neoplasms can be distinguished from other diseases in 90% of cases. Among the conditions to be differentiated are chronic inflammatory or toxic states, such as meningeal tuberculosis or parasitic granuloma; congenital anomalies with hydrocephalus; and trauma causing subdural hematoma.

To determine the site and nature of brain tumor in a child may be difficult or impossible. With no focal signs, the lesion is probably in the posterior fossa or third ventricle. In the latter site, malignant growth is practically certain.

In doubtful cases, the posterior fossa is explored. If no tumor is found, a Torkildsen ventriculostomy is done, and roentgen therapy is applied.

**B**ACTERIAL PNEUMONIA in children usually subsides quickly with Chloromycetin therapy. Fever of pneumococcal, streptococcal, or unclassified type commonly disappears in three days or less, and pulmonary lesions resolve in about six days, find Adrian Recinos, Jr., M.D., and associates of the Children's Hospital, Washington, D.C. When from 50 to 220 mg. of Chloromycetin per kilogram of body weight was given daily by mouth in capsules or syrup to 32 patients, all but 1 recovered promptly. The only unfavorable reaction was slight temporary depression of leukocytes

*New England J. Med.* 241:733-737, 1949.



## Hyperthyroidism after Sixty

ELMER C. BARTELS, M.D., AND J. W. KINGSLEY, JR., M.D.\*

*Lahey Clinic, Boston*

SIGNS and symptoms indicative of hyperthyroidism in elderly patients are often insignificant or are interpreted as the effects of old age. Yet 12% of all cases of the disease occur after the age of sixty years. As in the earlier decades, females predominate, the ratio being 4 women to 1 man.

Besides an enlarged thyroid, the most distinctive sign of hyperthyroidism in the aged is increased warmth of hands and body, observe Elmer C. Bartels, M.D., and J. W. Kingsley, Jr., M.D.

Hyperthyroidism due to a toxic adenomatous goiter occurs more frequently than primary hyperthyroidism. Although almost all such persons gradually lose weight, the chief symptom noted by the patient is usually general weakness or debility. Exertional dyspnea and palpitation are common.

Patients with primary hyperthyroidism usually first notice a rapid weight loss. Muscular weakness and cardiac palpitation are also often described.

Nervousness and emotionalism appear in but a small percentage of cases, whether the disease is primary or secondary. Usually the appetite is unchanged but anorexia is more common than increased appetite. Extended fingers often display the typical fine tremor.

A nodular goiter is usually readily palpable, but with primary hyperthyroidism the gland is often only slightly enlarged.

Tachycardia is rare. A resting pulse rate in excess of 120 per minute is the exception in the elderly hyperthyroid patient. Auricular fibrillation occurs about one-third of the time with toxic nodular goiter. One-fifth of patients with primary hyperthyroidism have this arrhythmia.

Congestive heart failure is apparent in 14% of the cases of primary thyrotoxicosis, and in 19% of secondary hyperthyroidism.

Clinically, hyperthyroidism in the geriatric patient may be called apathetic. However, the elevation in the basal metabolic rate is similar in degree to that found in young patients. Over half of old thyrotoxic and 40% of secondary hyperthyroid patients have basal metabolic rates of 40 or above.

Therapy is begun with a thiourea compound; 600 mg. of thiouracil or 200 to 300 mg. of propylthiouracil is usually given daily. This dosage causes the basal metabolism rate to decrease about 1% each day in primary hyperthyroidism. The return to euthyroidism is slower with toxic adenoma.

In many cases, in order to help prepare the patient for surgery, propylthiouracil therapy is continued for

\* Hyperthyroidism in patients over sixty. *Geriatrics* 4:535-540, 1949.



some time after the basal metabolism rate has regressed.

With a diffuse toxic gland, Lugol's solution is given for three weeks before surgery. For auricular fibrillation, preoperative digitalization is indicated.

A one-stage subtotal thyroidectomy is performed. With proper medical

preparation, the postoperative morbidity is slight. Of 123 patients thus treated only 1 died. Most patients are able to leave the hospital six days after operation.

The usual complications of thyroidectomy must be watched for. Postoperative myxedema requires thyroid extract.

## Functional Sterility and Amenorrhea

RITA S. FINKLER, M.D.\*

**I**RRADIATION may be preferable to hormone therapy for pituitary-ovarian hypofunction.

Therapeutic effects in functional sterility and amenorrhea are achieved much more quickly and slightly more often by roentgen rays than by gonadotropins, explains Rita S. Finkler, M.D.

Endocrine deficiency was observed in 190 married or single women at the Beth Israel Hospital, Newark, N. J. Among the group desiring children, conception took place after irradiation in 35% of cases and after medication in 34%. Menstruation was restored in 46% of patients given irradiation and in 41% of those given hormones.

The endocrine status is determined by endometrial biopsy, vaginal smears, basal temperatures, and urinary gonadotropin titers.

Equine or anterior pituitary gonadotropins are injected four to six times daily or on alternate days in doses of 200 to 500 units in the first half of the cycle. Chorionic gonadotropins may be given in the second half or combined with anterior pituitary in the first. Hormones are sometimes continued six or eight months.

Since irradiation is not advisable after conception, courses for menstruating women are begun just after the last period, and intercourse is banned during treatment. With amenorrhea, a preliminary Friedman test is done.

Therapy requires 200 kilovolts, with 0.6 mm. copper and 1 mm. aluminum filtration at 50-cm. target-skin distance. The half value layer is 1 mm. copper. Ovarian fields are 8 by 10 cm. and pituitary 6 by 8 cm. In most cases 4 treatments are given in two weeks, and each organ receives 80 r.

\* Evaluation of hormonal and radiation therapy in 190 cases of functional sterility and secondary amenorrhea. *Am. J. Obst. & Gynec.* 58:359-364, 1949.

## Causes and Effects of Eclampsia

CHARLES I. BRYANS, JR., M.D., AND RICHARD TORPIN, M.D.\*

*University of Georgia, Augusta*

SOME constant environmental or dietary condition is more likely than inherent physiologic weakness to cause eclampsia of pregnancy. Because the etiologic agent is unaltered, toxemia is apt to reappear with subsequent pregnancies and to increase the chances of abortion or stillbirth.

Charles I. Bryans, Jr., M.D., and Richard Torpin, M.D., believe that eclampsia is neither a manifestation nor a cause of chronic nephritis or hypertensive cardiovascular disease, although either of these conditions may precede an attack of eclampsia and possibly render the patient susceptible to toxemia.

For an average of twelve years, data was collected on 243 women with eclampsia during pregnancy; 138 were white and 105 Negro. After the eclamptic gestation, 188 of the women had a total of 565 pregnancies.

Over half the patients were toxemic at least once again.

Nearly one-fourth of conceptions occurring after the first toxic pregnancy resulted in abortion or stillbirth. Apparently, fetal mortality for women subject to eclampsia is almost twice the rate for the general population.

True eclampsia was repeated in almost 1 of 20 cases, or at least 7 times as often as is expected in usual

birth statistics even by the most conservative estimate. Before the original severe attack, however, toxemia was no more frequent than for all parturients.

Of the 243 women, 27 died within one to twenty-eight years. Approximately 14% of these succumbed to eclampsia, 15% to chronic glomerulonephritis, and 18% to cardiovascular disease.

The incidence of hypertension was not remarkably high, although younger women of the eclamptic group were more often involved than would be expected in an unselected sample. Both before and after the original severe attack of toxemia, the hypertensive group actually had fewer instances of toxic pregnancy, abortion, and stillbirth than those with normal postpartum blood pressure.

Thus eclampsia does not cause hypertensive cardiovascular disease. However, the toxic state perhaps aggravates an established vascular disorder, so that cardiovascular symptoms appear earlier than usual.

Convulsions nearly always result from long-neglected toxemia, hence conclusions drawn for eclampsia probably hold for preliminary phases. No permanent vascular lesions may then be expected from either pre-eclamptic or eclamptic toxemia. If hypertension is observed after a pregnancy with nonconvulsive involve-

\* A follow-up study of two hundred forty-three cases of eclampsia for an average of twelve years. *Am. J. Obst. & Gynec.* 58:1034-1065, 1949.

ment, the supposedly preeclamptic condition may have been essential hypertension from the start.

However, true preeclamptic toxemia can and does occur after eclampsia. Women with normal blood pressure after the episode subsequently had about 3 pregnancies, of which

more than one-third were toxemic. In a lengthy period of observation, any latent or potential hypertensive cardiovascular condition would certainly have become evident, and the toxic state must therefore result from some other cause than essential hypertension.

## A Nonmoving Ligature-Holding Bobbin

JOHN DEVINE, M.D.\*

THE principal of the fisherman's spinning reel, in which the line is peeled from the end of a stationary spindle, has been adapted to a ligature-holding bobbin by John Devine, M.D., of Melbourne, Australia.

The bobbin has no moving parts. Resistance, as the suture material is drawn out, is almost imperceptible, and snarling and jamming are avoided.

The bobbin, made of solid chromium-plated brass, consists of a core or spool and an outer casing. Stainless steel threaded tubes of any shape or length may be fitted to the end of the outer casing as a ligature carrier.

When necessary or desirable to tie at a depth with the suture material at the ligature carrier point, as in a tonsillectomy, braking pressure is applied by pushing the inner bobbin from the rear, so that the suture material is held between the inner bobbin and the edges of the end hole in the outer casing.

For underrunning vessels, as in an appendectomy or gastrectomy, the bobbin can be used alone or with a 3-in. slightly curved ligature-carrier, which can be braked by the thumb while the bobbin is held in the hand.

When ties are to be used in two places and the vessel cut between, as in saphenous vein ligations, different colored suture materials, wound together, come out of 2 separate holes in a short, curved ligature passer. Thus, when the vessel is underrun, two ligatures are placed at once and the variation of colors helps to prevent their being crossed.


\* A ligature-holding bobbin without moving parts. M. J. Australia 2:675-676, 1949.




## Hemorrhoidectomy

F. M. AL AKL, M.D.

Kings County Hospital, New York




KEEP THIS PICTURE IN MIND




1. Place patient face down; adjust table to lower trunk and elevate buttocks. Fix gluteal fold open with T adhesive straps.




2. Paint and drape field. Lubricate gloved finger; palpate anal canal and rectal wall.




3. Insert bivalve speculum and gently dilate anorectal ring. Rotate speculum and inspect anal canal and rectum.



4. Insert small sponge on holder into rectal ampulla and rotate; gently pull out on holder, exposing extruded hemorrhoidal masses and redundant mucosa.




5. Apply triangular clamp radially to crest of each presenting primary hemorrhoid, and to secondary masses when present. Withdraw spongystick.



6. Lift triangular clamp and make acute V-shaped incision from outer angle of hemorrhoidal ridge to beyond anorectal line.




7. Reflect anoderm from mass as assistant opens V with fingers. Clamp bleeders.




8. Continue dissection with scissors, freeing undersurface of hemorrhoidal mass from subcutaneous band of external sphincter.



9. Transfix pedicle containing hemorrhoidal vessels; ligate and excise mass distal to ligature. Tie bleeders and cut ligature.




10. When mucosa is redundant, clamp mass radially after hemorrhoid is dissected free.



11. Pass suture ligature below tip of clamp transfixing edge of mucosa containing vessels. Tie ligature and excise mass.







12. Whip rest of suture loosely over clamp. Open and withdraw clamp.



13. Pull on suture and tie ends together; inspect for bleeding, then cut ligature.



14. If suture line is long or retracts too far into rectum, pull taut and anchor beyond sphincter band and tie.



15. Repeat procedure on remaining hemorrhoids, leaving intact strips of anoderm between incisions.

#### NOTES

With regional anesthesia, the jack-knife position affords good exposure and an accessible operative field. When general anesthesia is administered, the lithotomy position is easier for the patient.

Neat incisions and careful dissec-

tion of hemorrhoidal masses without crushing or macerating anoderm reduce patient's discomfort. Oily anesthetic injections and rectal tubes are unnecessary when the dissection is clean and the edges of the anoderm undamaged.



## Blood Pressure and Surgical Position

L. H. PETERSON, KENNETH F. EATHER, M.D.,  
AND ROBERT D. DRIPPS, M.D.\*

*University of Pennsylvania, Philadelphia*

A NEW method of recording blood pressure during operation instantly shows the effect of postural change.

Intra-arterial and pulse pressures are traced from beat to beat by the compact mobile ink-writing manometer and a small plastic catheter of special design. The pulse wave shows whether a sudden drop results from splanchnic pooling or decreased peripheral resistance.

Abrupt hypotension has often been noted during operations, state L. H. Peterson, Kenneth F. Eather, M.D., and Robert D. Dripps, M.D., when the patient's legs are moved from the lithotomy to the horizontal position.

The flexible catheter is made of synthetic polyvinyl resin tubing. A section is heated in an oven for seventy-two hours at 110° C., then drawn out to an inside diameter of about 0.2 mm.

A 1-cc. tuberculin syringe with a 22-gauge needle and hollow steel plunger is used to insert the catheter. One end of the plunger has a nipple for attachment of the catheter, and the other end fits a capacitance manometer with small volume displacement.

The syringe barrel is filled with sterile physiologic saline solution, the

needle thrust into the blood vessel, the plunger pushed down to inject the catheter, and the needle then withdrawn. The catheter can be left in the artery for hours without harm.

The manometric system is previously flushed with anticoagulant solution and during use receives a small amount from a pressure reservoir through a side tube and needle valve.

Since the recording unit is connected to the patient only by 3 small flexible leads, tracings are made with little disturbance of surgical routine. The apparatus can be moved 20 ft. away, shifted from one subject to another by change of catheter, and operated by a single person.

Even among healthy individuals, a tilt from supine to erect position produces circulatory collapse in 8 to 10% of cases. If vessels in skin and muscle are dilated by sympathetic blockade or resection, heat, or other means, the reaction is more severe. After intramuscular administration of 15 mg. of morphine sulfate, tilting may cause profound hypotension in about half of patients.

The patient under sedation or deep anesthesia during surgery is thus susceptible to postural change. After a vaginal procedure, perineal prostatectomy, or other operation, lowering of raised legs may reduce

\* Postural changes in the circulation of surgical patients as studied by a new method for recording the arterial blood pressure and pressure pulse. *Ann. Surg.* 131:25-50, 1950.

systolic blood pressure as much as 60 mm. of mercury. A sharp decrease is likely after excessive hemorrhage or prolonged elevation causing anoxic damage to blood vessels.

To correct hypotension, legs may be raised at the hip without moving the trunk, held up about two minutes, lowered, and if necessary lifted once or twice more at short intervals. For failure due to peripheral vasodilatation, this maneuver is physiologic and acts more rapidly; it succeeds more often than pressor drugs. Circulation is improved without the respiratory embarrassment of the Trendelenburg position.

If the fall in blood pressure results from collection of blood in the viscera and diminished stroke volume, both postural change and pressor drugs may be useless.

To determine the nature of circulatory collapse, the pulse pressure contour should be analyzed. With lowered peripheral resistance and pooling in the extremities, the height of systolic rise is practically unaltered, but the rate of diastolic fall is more rapid than before blood pressure dropped. Peripheral constriction and lessened stroke volume reduce the systolic rise and broaden the peak.

**SERUM ACID PHOSPHATASE** should not be determined for twenty-four hours after prostatic massage. Within the first hour after massage, values may increase to those for widespread prostatic cancer, report Ernest Hock, M.D., and Roland N. Tessier, M.D., of the Wilson Memorial Hospital, Johnson City, N. Y. Effects of massage were noted in 20 cases. With good renal function the acid phosphatase concentration was highest within an hour after massage, occasionally elevated for about five hours, and at or slightly below the original estimate in twenty-four hours. When kidneys are impaired the rise may persist for two days.

*J. Urol.* 62:488-491, 1949.

**BACILLARY URINARY INFECTIONS** are usually suppressed or reduced by Chloromycetin. From 1 to 1.5 gm. per day is given in divided doses of 0.25 to 0.5 gm. at intervals of six to eight hours. With severe involvement, the first dose is 1 gm. Since coccal organisms are often present, penicillin, sulfonamide, or both may be required. George E. Chittenden, M.D., and associates of Charles Godwin Jennings Hospital, Detroit, treated 50 patients with Chloromycetin, for most of whom other antibiotics had been unsuccessful. The majority of patients had calculi, enlarged prostate, or other lesions. Symptoms disappeared in 82% of cases and urine became sterile in 38%. The only toxic reaction observed from the antibiotic was slight vertigo in 1 instance.

*J. Urol.* 62:771-790, 1949.

## **starting point for all feeding problems...**


Almost every type of infant formula can be simply made when Dryco is used as a starting point. Dryco is a versatile formulation which serves as a basis for designing special diets for healthy infants or during illness and convalescence.

In itself Dryco assures ample protein intake for optimal body growth and tissue development. Dryco provides less fat, thus minimizing the danger of digestive disturbances. Carbohydrates may be added when indicated, and the fat content may be modified by adding milk (fresh, evaporated or dried) to the formula.

*Moreover...* Dryco's special drying process makes it more easily digested than the fresh milk from which it is made.

Dryco supplies more minerals, particularly more calcium, than a corresponding formula of whole milk.

Dryco contains those vitamins in fluid milk which are essential for infant health and growth. Only vitamin C need be added.



Dryco is made from spray-dried, pasteurized, superior quality whole milk and skim milk, providing 2500 U.S.P. units of vitamin A and 400 U.S.P. units of vitamin D per reconstituted quart.

Each tablespoonful supplies 31½ calories. Readily reconstituted in cold or warm water. Available at pharmacies in 1 and 2½ lb. cans.

**Dryco**® *a versatile base for "Custom" formulation*

The Prescription Products Division, The Borden Company  
350 Madison Avenue, New York 17



# Medical Forum

*Discussion of articles published in MODERN MEDICINE is always welcome. Address all communications to The Editors of MODERN MEDICINE, 84 South 10th St., Minneapolis 3, Minn.*

## Abdominal Paracentesis\*

TO THE EDITORS: The continuous paracentesis by the electrosurgical method of Dr. William James Moore is novel and interesting. He has indicated that the advantages of his technic are prolonged drainage and less risk of wound complications. Critical evaluation of this procedure should be reserved if one has not had actual experience in its use. However, there seem to be some disadvantages in his method which should be mentioned.

Insertion of the electrocoagulating trocar leaves a zone of necrotic tissue through all layers of the abdominal wall. Charred, devitalized tissue is an inviting culture medium for bacterial invasion, either from within or without the abdomen. After the slough of electrocoagulation has been discharged or absorbed, the burned tissue tends to heal slowly with thick scar formation.

For those less skilled than Dr. Moore, caution should be taken not to damage intestine with the electrocoagulating trocar as it is being inserted. If the thin intestinal wall contacts the active electrode a fatal peritonitis might result from leakage at the site of injury.

Experience in the surgical research

\*MODERN MEDICINE, Nov. 1, 1949, p. 58.

laboratory and with clinical problems of abdominal drainage has shown that organic substances, such as rubber, gauze, or plastic materials, produce a marked fibroplastic response in the peritoneal cavity which results in a relatively rapid walling-off of such drains. This interdicts the use of drains made of organic substances when continuous drainage is required. Conversely, materials of inorganic structure, such as glass or stainless steel, produce minimal reaction when placed in the peritoneal cavity. Efficient prolonged abdominal drainage by gravity or suction can be more satisfactorily obtained when the intraabdominal portion of the drain is constructed from an inorganic substance.

DANIEL J. PRESTON, M.D.

Wilmington

► TO THE EDITORS: There is no doubt that the many problems associated with abdominal paracentesis are so distressing that any contribution to the eradication of these problems is well received.

As for the advantages of the electrosurgical technic, I see but two assets. These are the prevention of implanting malignant cells in the wound and, secondly, the sterilization

of the tract without the dissemination of pathogenic bacteria. Dr. William James Moore maintains that tuberculosis may be treated well by this method. Tuberculous peritonitis often responds simply to pneumoperitoneum following the routine type of paracentesis. Apparently the author's method does not allow air to enter the peritoneal cavity. Thus a beneficial introduction of air is not possible in tuberculous ascites.

BERNARD J. FICARRA, M.D.

Brooklyn

### Temperature Cycle in Obstetrics and Gynecology\*

TO THE EDITORS: Dr. Melvin B. Sinykin has presented an excellent résumé of the relationship between ovulation and rise in basal temperature of the body.

As he points out, the body temperature in most menstruating women follows a cyclic pattern and the temperature curve often reflects certain phases of the menstrual cycle. Such cyclic variations are not seen in women who are not truly menstruating, that is, who are not ovulating. However, as in all matters in medicine, the rule does not always hold and these cyclic variations are not always clear, even in ovulating women.

For women who can practice rhythm contraception, basal temperature makes an excellent guide. However, every woman cannot follow this method; some conceive even when intercourse takes place only during menstruation, although such conception would seem impossible in view

\*MODERN MEDICINE, Apr. 1, 1949, p. 64.

of our present beliefs on the time of ovulation and the length of life of the spermatozoon.

P. A. MCLEOD, M.D.

Kingston, Ont.

### Amyotrophic Lateral Sclerosis

TO THE EDITORS: As a result of the publication of my statement in your Medical Forum that adrenal cortex therapy appeared to influence favorably the course of amyotrophic lateral sclerosis (*Modern Medicine*, July 15, 1949, p. 72), I have had numerous written requests for details and, more recently, privately communicated confirmation of the myodynamic "lift" afforded some patients. I also mentioned a more insensible sustaining effect which I believed to become overt only after prolonged therapy.

I am impelled to elaborate somewhat on my present treatment schedule at this time, first, because the written requests have been for further information and, second, because a definitive paper, now in preparation, will probably not see publication for many months. There is the further impetus of a very recent verbal report at a Chicago Symposium that amyotrophic lateral sclerosis was one of the diseases *not* favorably influenced by the administration of adrenocorticotrophic hormone, ACTH.

In view of my own present conviction that adrenocortical therapy justifies its onerous expense to patients with this disease, I am anxious that the single report of the failure of adrenocorticotrophic therapy be not misconstrued. There may be a difference and an explanation of the difference.

## MEDICAL FORUM

With regard to my treatment schedule, it consists of the individualized administration of from 3 to 15 cc. of whole cortex extract (aqueous) daily, or in divided doses twice daily. The dose which reduces fasciculations, spasticity, and ankle clonus to minimums is found and continued indefinitely.

Occasional trials at dosage reduction were not well borne by my patients, but they responded anew to resumption of full dosage. These trials had been instituted to alleviate possible pituitary adrenocorticotrophic depression; they inevitably led to a degree of relapse that made apparent, by contrast, the previous gradual improvement of the patients during periods of relentless cortical extract administration. I no longer worry about the pituitary, and though I continue B<sub>12</sub>, a high-fat diet, testosterone, and progesterone (in some instances) as a matter of orthodoxy, I am reasonably sure that adrenal cortex is the prime factor in causing the arrest.

Evaluation of results should be made only after some months of continued therapy. In such an evaluation, no reasonable person would anticipate any extensive restitution of function of markedly demyelinated tracts. Any mode of treatment must be evaluated in light of the physiologic dogma that there can be no regeneration, anatomically, within the central nervous system proper. Long-standing cases of amyotrophic lateral sclerosis are poor objects for reexamination of the facets of this dogma, but I have seen indubitable evidence of definite, if minimal, return of volitional activity to certain long-dormant muscle groups.

Amyotrophic lateral sclerosis is probably an atopic response to allergenic cholinergesis, with or without psychogenic cholinergesis, in which the pyramidal pathways and their connections react as the atopic shock organ. As in all atopic reactions, two independent modalities of compensation are called into play. One is the Selye corticotrophic-cortical "alarm" mechanism and the other is the erythropoietic cholinesterase mechanism.

The latter modality alone fails in such a condition as acute leukemia, which is also the result of an atopic reaction. Both modalities may fail in irreversible surgical shock. In amyotrophic lateral sclerosis there is usually enhanced erythropoiesis and an elevated blood cholinesterase so that, on this premise, compensation failure must lie in the corticotrophic-corticosteroid mechanism. Extension of the premise furnishes the rationale of adrenal cortex therapy.

ROBERT D. BARNARD, M.D.

New York City

### Diet in Diabetes\*

TO THE EDITORS: Dr. Lester J. Palmer presents a dietetic regimen for the diabetic, with a sample menu containing carbohydrate 165, protein 80, and fat 90 gm. His views are in agreement with the majority of authoritative opinions—although there are notable dissenters.

It has been the custom to provide 1 gm. of protein per kilogram of body weight. There is increasing evidence that adequate stores of body

\*MODERN MEDICINE, June 1, 1949, p. 57.



protein are helpful in resisting infection and in prevention of fatty livers and, possibly, of retinitis. Even higher values than the increased protein allowance suggested by Dr. Palmer are therefore advocated by some authorities.

The optimum carbohydrate in the diet is still a subject of debate, ranging from the unlimited choice of the "free-dieters" to that of strict limitation. Diets containing 150 to 200 gm. are most popular. Smaller amounts are disliked by patients, while higher starch values make attainment of near normal blood sugars more difficult. The desirability of the latter objective in itself has been much debated.

Recent experimental evidence appears to support the view that hyperglycemia is damaging to the pancreas. However, the ideal of maintaining rigid control of blood sugar levels will in many cases evoke hypoglycemic reactions. The damage which these reactions are believed to cause to the central nervous system would outweigh any benefits gained by sparing the pancreas.

Having provided for the protein and carbohydrate intake, the remaining calories required will be made up of fat. However, there are some who believe that fat should be fixed at a low level, with consequent increase in carbohydrate, on the theory that the incidence of hypercholesterolemia and atherosclerosis can be reduced thereby. These diets are somewhat unpalatable and it is also possible that deficiency in essential food factors may occur. Moderate fat allowance, such as recommended by Dr. Palmer, is therefore most com-

monly in use. The problem of the relation of diet to arteriosclerosis is not yet settled, and an open mind should be maintained on this question.

B. H. LYONS, M.D.

Winnipeg

► TO THE EDITORS: The diet presented by Dr. Lester J. Palmer is for the uncomplicated diabetic. It is liberal with respect to carbohydrates and protein. It contains more fat than the high-carbohydrate low-calorie diet, but this circumstance is, to some extent, counterbalanced by the fact that the caloric content is regulated by the energy requirements of the individual.

Dr. Palmer also stresses *variety* of food materials. The diet thus meets the vitamin and mineral requirements without subjecting the diabetic to the inconvenience and expense of special preparations of these essential food elements. The diet, therefore, conforms to well-established physiologic principles. Conforming also to well-established practice is the insistence upon sugar-free urine and a blood sugar level approximating normal.

Compared with this method of treatment is the increasing use of the "free" diet combined with insulin, with little or no attention to blood and urine sugar—a practice which is not only uneconomical but inevitably harmful if continued for long. With the free diet, not only do many more diabetics require insulin than with a diet such as Dr. Palmer's, but the average dose is appreciably larger.

## MEDICAL FORUM

Clinical experiences and animal experimentation also clearly show that persistent hyperglycemia, which is almost inevitable with the free diet, results in deterioration of the islands of Langerhans and also of the liver, as shown by liver function tests, enlargement of the liver in children, and fatty infiltration at autopsies. Dr. Palmer's criteria of the control of diabetes are therefore particularly gratifying.

An additional consideration with the free diet is that the increasing diabetic population will inevitably aggravate the problem of insulin supply, unless, in the meantime, new sources of insulin are discovered.

I. M. RABINOWITCH, M.D.

Montreal

### Culdoscope in Gynecology

TO THE EDITORS: I feel that Drs. Richard B. Dunn and Donald C. Schweizer have covered the subject of the culdoscope very well and that this method of diagnosis has a very definite place as a diagnostic procedure in gynecology. I have been interested in the use of the culdoscope ever since Dr. Decker first described its use. Through his efforts in the utilization of the knee-chest position he has made peritoneoscopy a valuable diagnostic adjunct.

We have not considered general anesthesia necessary for culdoscopic examinations and have done many satisfactorily by the local injection of 1% novocain into the vagina after the patient has been well sedated. However, lately, we have resorted to the use of saddle-block type of anesthesia with excellent anesthetic ef-

\*MODERN MEDICINE, Jan. 1, 1950, p. 67.

fect. There is definitely more difficulty in having the patient retain the knee-chest position when saddle-block is used than when local anesthesia is given.

This culdoscopic procedure has a definite place in obviating many laparotomies that are done simply for exploratory reasons. The surgical risk with culdoscopy is much less than with laparotomy. The entire pelvis can be viewed quite satisfactorily and pathologic processes diagnosed, with sufficient training.

As the authors state, this procedure is also very valuable in the diagnosis of sterility, inasmuch as the patency of the tubes can be discerned by direct vision when the culdoscopy is done.

The contraindications of this procedure are few: any fixed mass in the cul-de-sac, and cardiac conditions that preclude the patient's assuming the knee-chest position.

HARLEY E. ANDERSON, M.D.

Omaha

► TO THE EDITORS: We have been using the culdoscope as a diagnostic aid for several years and think it is a great help in many cases.

We have made a light, wooden, padded board shaped like an inverted "U," which rests on the operating table to hold the patient in knee-chest position during the procedure. It has been a great asset, especially if the patient has received pentothal.

Our main difficulties with the procedure occur when adhesions obscure the field or with an adherent retroverted uterus.

WILLIAM BUSTER MCGEE, M.D.

San Diego

# What Can Be Done for the Hay Fever Patient?

TO THE EDITORS: The excellent paper by Dr. Albert V. Stoesser refers especially to the treatment of hay fever in children, but most of the statements made are equally applicable to adults.

I heartily agree with the author that hay fever patients should be regarded as perennial problems. If we could persuade the patient that his treatment is a matter of years instead of the particular year in which he consults the physician, the results would be very much more satisfactory.

I was pleased to see that the use of drops in the nose was not advocated except for a very short period of time. Much harm has been done by instillation treatment.

The author's comments on antihistamine therapy are quite helpful and the classification of the various antihistaminic drugs is clarifying.

Of course, the injection treatment of pollen extracts is rightly considered the most important part of the treatment. Careful reading of this section of the article is well worth while and I would especially emphasize the statement made that "approximately 60% of the failures were due to lack of cooperation on the part of the patient, the doctor's office, or both in following closely the dosage schedule, to absence of information concerning other allergies of the hay fever sufferer, and to full disregard of special rules of conduct during the hay fever season."

HERBERT K. DETWEILER, M.D.

Toronto

\*MODERN MEDICINE, Aug. 1, 1949, p. 47.

► TO THE EDITORS: I was interested in reading Dr. Stoesser's article. Most of his hay fever cases have been in children and he has given an up-to-date review of the best methods of diagnosis and treatment.

As he points out, many of these children do not present the usual symptoms of rhinitis and corneal injection, but may have a continual irritating cough or a "summer cold" which lasts for some weeks or even a few months, then clears up suddenly when the pollen count drops. The symptoms will be repeated the next summer and this time the physician should realize that the inhalation of grass or ragweed pollen is the real cause of the patient's symptoms.

Dr. Stoesser rightly stresses the fact that while the newer antihistamine drugs have given temporary relief to many patients, they will not supplant the more reliable method of treating these patients by pollen injections. If the diagnosis is correctly made by skin tests, using the pollens or molds to which the patient is sensitive, and the patient is treated with these particular allergens, the results of treatment are usually very satisfactory, particularly if the patient is a child and has not had symptoms for more than a few years.

I do not altogether agree that the preseasonal method of treatment is preferable to the perennial method of giving doses of the specific pollen at long intervals throughout the year. We believe that a higher degree of desensitization may be achieved by this latter method.

J. R. ROSS, M.D.

Toronto

# Diagnostix

*Here are diagnostic challenges presented as they confront the consultant from the first clue to the pathologic report. Diagnosis from the Clue requires unusual acumen and luck; from Part II, perspicacity; from Part III, discernment.*

## Case MM-162

### THE CLUE

ATTENDING M.D.: I remember your telling me not long ago that you were especially interested in hemolytic anemias.

VISITING M.D.: Yes, and I still am.

ATTENDING M.D.: Good. Then you'll want to see a two-year-old child that has just been admitted with



Lederer's anemia. We presented him at the last weekly clinic. He has been in the hospital only twelve hours and is receiving transfusions.

VISITING M.D.: Lederer's anemia is one of an extremely interesting group of blood diseases—the so-called acute hemolytic anemias of unknown origin. The disease is usually attributed to infection, you know. Did the

child have a febrile illness, leukocytosis, lethargy, and gastrointestinal symptoms?

ATTENDING M.D.: Well, yes, lethargy and diarrhea and he is feverish. The leukocyte count is 20,200, with 60% neutrophils, 36% lymphocytes, and 4% eosinophils. Red cell count 2,000,000; hemoglobin 5 gm. per 100 cc., and hematocrit 20 volumes per cent. Smear showed anisocytosis, fragmentation, and polychromatophilia. Urine was port wine in color, and the benzidine test for hemoglobin was strongly positive.

VISITING M.D.: What caused the anemia? How acute is the illness?

ATTENDING M.D.: I do not know. The child was in good physical health until three days ago when he started passing six to eight loose stools a day and wine-colored urine. According to his parents, he had received no medication and had not been exposed to any poisons or chemicals.

### PART II

VISITING M.D.: Porphyrins negative? Other studies?

ATTENDING M.D.: No porphyrins in the urine. The reticulocyte count is 8%. Sternal aspiration shows nothing diagnostic. Serologic test for syphilis not reported. No icterus.

VISITING M.D.: Acute hemolytic anemia

ANNOUNCING  
A NEW MEMBER  
OF THE  
FURACIN FAMILY



**FURACIN**  
**ANHYDROUS EAR SOLUTION**

In the treatment of bacterial OTITIS media et externa, over 200 reported cases attest that Furacin is a highly effective adjunct.\* Many cases of chronic otitis responded which had proven refractory to other medicaments. Among the pathogens isolated were *Escherichia coli*, *Proteus vulgaris*, *Pseudomonas* species, staphylococci, streptococci and diphtheroids. Furacin Anhydrous Ear Solution contains Furacin® 0.2 per cent, brand of nitrofurazone N.N.R. in an anhydrous, hygroscopic, water-soluble liquid: polyethylene glycol. It is indicated for topical treatment of bacterial otitis media et externa. *Literature on request.*

EATON LABORATORIES, INC., NORWICH, N. Y.

\*Anderson, J. and Steele, C.: *Use of Nitrofurazone Therapy in External Otitis*, Laryngoscope 58:1279, 1948 • Douglass, C.: *The Use of Furacin in the Treatment of Aural Infections*, Laryngoscope 58:1274, 1948 • Reardon, H.: *Unpublished results.*



## DIAGNOSTIX

and hemoglobinuria of unknown etiology. . . . Let me talk to the father and mother. (*A long interview ensues in the office, the consultant then comes out*) Let me see a sample of the stool and urine. (*Smells the specimens*) Send a specimen of urine to the toxicology department.

### PART III


VISITING M.D.: (*Examining patient*) Heart, lungs, abdomen negative. Fundus—roughly negative. Deep reflexes—normal, nothing objective except temperature of  $100.2^{\circ}$  rec-

tally. The father told me that the boy had been chewing moth balls.

### PART IV

ATTENDING M.D.: (*Two days later*) The toxicologist reported that the first urine specimen contained naphthol; yesterday's contained only a trace; and today's, none.

VISITING M.D.: The moth balls that the parents brought in contained naphthalene. Moth balls are far from harmless, yet children occasionally suck them. I have seen a similar case. Well, we've lost our case of Lederer's syndrome.





*National Synthetics*

*takes pleasure in inviting*

*you to try*

*Monophen Tablets*

*the new, better-tolerated,*

*versatile medium*

*for*

*oral cholecystography*

*R. S. V. P.*

*National Synthetics, Inc.  
270 Lafayette Street  
New York 12, N. Y.*

*Write for Clinical Samples and Literature*

**NATIONAL SYNTHETICS, INC.**

270 LAFAYETTE STREET, NEW YORK 17, N. Y.



# *A Valuable Adjunct in* **INFANT and CHILD FEEDING**



## *Unflavored Gelatine as an Aid to Nutrition and Digestion*

The unique qualities of Knox Unflavored Gelatine make it especially useful in the dietary of infants and children.

In certain cases where difficulty is encountered in the digestion of a milk formula the addition of 2 percent of Knox Gelatine improves digestibility. It is also useful in increasing the protein content of the formula.

Older children benefit from the recognized nutritional benefits and added appetite appeal of gel-cookery dishes made with Knox. Unlike ready-flavored gelatin dessert powders with their high sugar and acid content, Knox is all protein with no sugar content.

**FREE** DIETARY GUIDE BOOK IN  
INFANT AND CHILD FEEDING

"Knox Gelatine in Infant and Child Feeding" contains detailed infor-

mation compiled from experience, plus an assortment of carefully selected recipes. Copies are available to you upon request. Write Knox Gelatine, Dept. R-25, Johnstown, N. Y.

Available at  
grocery stores  
in convenient  
4-envelope and  
38-envelope  
packages.



# **KNOX**

Gelatine U. S. P.

ALL PROTEIN

NO SUGAR

**three  
essentials**...in the effective treatment of biliary dysfunction  
are combined in Caroid and Bile Salts Tablets.

They afford a desirable threefold action as—



**K**  
**choleric**...to produce increased bile flow

**Q**  
**digestant**...to assist digestion

**J**  
**laxative**...to induce peristaltic action and  
help reestablish normal function.

Caroid and Bile Salts Tablets have been a prescription  
favorite for many years as an effective aid in the treatment of  
biliary stasis, cholecystitis, constipation of biliary origin.  
Available in bottles of 20, 50, 100, 500 and 1000.

tablets of... **Caroid and Bile Salts** with Phenolphthalein

American  
Ferment

trial supply on request

Company, inc. 1450 Broadway, New York 18, N. Y.

# Short Reports

## THERAPY

### Acute Gouty Arthritis

Colchicine administered with or shortly after ACTH apparently prevents recurrence of acute gouty arthritis. Although hormone treatment alone is effective in combating the disease, Dr. William Q. Wolfson and associates of Chicago find that the glycocorticoid deficiency after withdrawal of ACTH may precipitate a new attack. Addition of colchicine to ACTH dosage appears to overcome this tendency. No patient given the concurrent dosage has had even a minor recurrence of acute gouty arthritis within one month of therapy.

*Proc. Central Soc. Clin. Research 22:93, 1949.*

## NEUROSURGERY

### Operation for Tachycardia

After complete sympathetic motor denervation, resting pulse rates are slowed, particularly in a patient with rapid rates. This procedure may be effective in hypertensive individuals with unusual degrees of tachycardia. By removing the second to fifth thoracic sympathetic ganglia on both sides, Dr. R. H. Smithwick of Harvard University, Boston, and associates noted reduction of basal pulse, acceleration after exercise, and no untoward effects. A few instances of exertional, emotional, or paroxysmal auricular tachycardia were also controlled by sympathectomy.

*Surgery 26:727-744, 1949.*

## ONCOLOGY

### Lung Cancer and Cigarettes

Smoke from cigarettes may be one of the causes for primary bronchogenic carcinoma. The disease, which was infrequent fifty years ago, is now one of the most common types of cancer. The increased incidence, suggests Dr. Evarts A. Graham of Washington University, St. Louis, is therefore attributable to some factor associated with modern living. The following observations indicate cigarette smoking as the responsible agent: [1] Statistical curves are similar for the increase both of bronchogenic carcinoma and of cigarette sales. [2] Lung cancer is rare in a patient who is not, or has not been at some time, an excessive cigarette smoker. [3] More men than women are affected by the disease, perhaps because women of cancer age seldom use cigarettes excessively. Use of tobacco in pipes and cigars apparently does not have the same etiologic relationship. The carcinogenic agent in cigarettes may be some factor used in curing the tobacco or in the paper or insecticides employed during plant growth.

*Inter-Am. Cong. Surg. 1949.*

## PERSONNEL

### Hawley to ACS


Dr. Paul R. Hawley, who recently resigned as executive officer of the Blue Cross and Blue Shield Commissions, has become director of the American College of Surgeons.

There is an important difference in Elastic Bandages!

# TENSOR\*

ELASTIC BANDAGE

is woven with *Live Rubber Thread!*



And *what* a difference that makes! Since TENSOR is woven with *live rubber thread*, it offers you these unequaled advantages over rubberless bandages.

## TENSOR:

- exerts uniform, controlled pressure without binding.
- has constant elasticity—you don't find it in rubberless bandages!
- stays put—needs no frequent readjustments—is comfortable to wear.
- is inconspicuous—women patients will wear it.
- can be laundered repeatedly without loss of elasticity.

**Ideal For Every Elastic Bandage Use!**

For your patients' support and comfort,  
prescribe Bauer & Black Elastic Supports

**BAUER & BLACK ELASTIC STOCKINGS**... 2-way stretch, easy to fit. Women appreciate their neutral and inconspicuous color.



**BAUER & BLACK SUSPENSORIES**... as a palliative and for treatment of diseases of the scrotum.



\*Reg. U. S. Pat. Off.

Products of

**(BAUER & BLACK)**

Division of The Kendall Company, 2500 S. Dearborn St., Chicago 16

**FIRST IN ELASTIC SUPPORTS**

## SHORT REPORTS

### NEUROLOGY

#### Epilepsy Sequela of Recurrent Malaria

Cerebral damage, manifested by convulsive seizures, may be caused by recurrent malaria. Since many soldiers discharged during World War II because of epilepsy were stationed in malarial areas when the seizures began, *Plasmodium* infection should be considered in the differential diagnosis of convulsive attacks, point out Dr. David R. Talbot and associates of Wadsworth General Hospital, Los Angeles. Electroencephalograms may prove useful in distinguishing the condition. If the epilepsy is a sequela of malaria, treatment of the malaria is of primary importance; anticonvulsive therapy is only secondary. Intensive malarial therapy may bring improvement before further damage appears.

J.A.M.A. 141:1130-1132, 1949.



"I must ask my patients to paint their house numbers on their roofs."

### ORTHOPEDICS

#### Molded Plastic Splints

A light, waterproof, and durable adjustable splint can be made of Celastic, a napped cotton material impregnated with pyrorylin. The method described by Beatrice F. Schulz of Washington University, St. Louis, is simple and inexpensive. The splint is cut from Celastic after being traced from a paper pattern. In forming a splint of more than two layers, the innermost or first and second layers are dipped in solvent and removed as soon as moist. The two layers are then laid together on a glass or rubberized silk surface and laminated by hand or roller pressure. When the fabric is no longer sticky and is dry enough to hold a slight curve, the laminated layers are shaped around the part to be splinted. The splint should remain in place until the material has set well enough to hold the proper contour when removed. An old prosthetic or discarded plastic shell can be used for rough shaping if the patient is not present. The Celastic must then dry for another hour before the finishing compound can be applied. During this time reinforcement layers and strap loops are added. About five or six coats of finishing compound should be applied to the splint either by dipping or by painting. Reshaping is possible after any of the applications. Because of the drying action of the solvent on the skin, rubber gloves should be worn while handling the moist Celastic, and the part to be splinted should be wrapped in non-porous material for initial fitting.

Physical Therap. Rev. 29:543-546, 1949.

# in choosing an oral xanthine preparation...

## *Check These Advantages*

- Low tendency to gastric irritation
- Oral dosage can be pushed to full therapeutic effect with minimal risk of unpleasant reactions
- Stability
- Ready absorption—brings about rapid clinical effect

BRAND OF THEOPHYLLINE SODIUM GLYCINATE

*Central*

*Raises the Xanthine Dosage Potential*

The oral administration of aminophylline and other theophylline preparations has heretofore been limited by their tendency to cause gastric irritation. Often the limit of gastric tolerance has determined the dose employed.<sup>1</sup> Theophylline-sodium glycinate has the advantage of being "less irritating to the gastric mucosa. It is thus tolerated orally in larger doses than are possible with other theophylline preparations, and it can be administered by mouth in liquid form as well as non-enteric-coated tablet form."<sup>2</sup>

### CHOOSE SYNOPHYLATE WHENEVER ORAL XANTHINES ARE INDICATED

**SUPPLIED: Tablets**—0.33 Gm. and 0.165 Gm., each 0.33-Gm. tablet equivalent to 0.16 Gm. Theophylline U.S.P. Bottles of 100, 500, and 1,000.

**Syrup**—Each teaspoonful (4 cc.) containing 0.33 Gm., equivalent to 0.16 Gm. Theophylline U.S.P. Bottles of 1 pt. and 1 gal. **Suppositories** (rectal)—Each suppository containing 0.78 Gm., equivalent to 0.39 Gm. Theophylline U.S.P. Cartons of 12, foil wrapped.

1. Goodman, L., and Gilman, A., *The Pharmacological Basis of Therapeutics* New York, The Macmillan Company, 1941; p. 281. 2. Council on Pharmacy and Chemistry, American Medical Association: *New and Nonofficial Remedies*, 1949. Philadelphia, J. B. Lippincott Company, 1949; p. 331.



THE CENTRAL PHARMACEUTICAL CO.

Products Born of Continuous Research

SEYMOUR • INDIANA



## SHORT REPORTS

### MENTAL HYGIENE

#### Child Mental Health

Enough knowledge on mental health has been gathered in recent years to permit a program of child care based on scientific principles. A national plan patterned after the one already in effect in Rochester, Minn., is suggested by Dr. Henry M. Helmholtz of the Mayo Clinic. The mental health experiment has been carried out for five years in the child health clinics of that city, which are used by virtually the entire population. Every child in the community is given a mental and emotional development test at the age of two-and-a-half years. When children's emotional disturbances are detected early enough, the large majority can be prevented from becoming personality difficulties. To make such a program effective, family physicians and school teachers as well as parents must learn the fundamentals of mental health.

### PUBLIC HEALTH

#### Red Cross Blood Centers

Thirty regional blood centers are now in operation in the United States, supplying blood for 1,550 hospitals in thirty-four states. Since the early part of 1948, the National Blood Program of the American Red Cross has provided more than 500,000 pints of blood for medical use. Taken from voluntary donors, the blood is furnished by the Red Cross centers to physicians and hospitals without charge. In addition to whole blood and plasma, valuable derivatives are distributed. The most important of these is immune serum globulin used as an immunizing or modifying agent

in measles. During the past year, the Red Cross distributed enough globulin for every case of measles reported in the United States. The derivatives were processed from surplus plasma returned by the Armed Forces after the last war. In another year, this supply will be depleted and globulin will have to be derived from blood received at regional centers.

### CARDIOLOGY

#### Chemical Pharmacology Section

The National Heart Institute has established a chemical pharmacology section under the direction of Dr. Bernard B. Brodie of New York University, New York City. Dr. Sidney Udenfriend of Washington University, St. Louis, will head the units within the section.


### CARDIOVASCULAR

#### Essential Hypertension

Veriloid is one of the most satisfactory of the extracts of *Veratrum viride* for treating patients with hypertension. Hypotensive action is similar to that of other *Veratrum* preparations and toxic effects fewer. The preparation can be administered for periods as long as five months. Dr. Robert W. Wilkins and associates of Boston University carefully adjust the dosage to the individual patient, increasing the amount slowly if necessary to lower the blood pressure, or decreasing the amount if the patient vomits or is nauseated. The initial dose, about 1 mg. of Veriloid four times a day after meals and at bedtime, may be slowly increased to about 2 mg. in three or four weeks.

*Proc. Soc. Exper. Biol. & Med.* 72:302-304, 1949.





In sickness or surgery, play safer with

## "Saturation Dosage"

of vitamins b and c

Depletion of the critical water-soluble B complex and C vitamins occurs so commonly in the presence of physical pathology, as to make a presumption of nutritive impairment<sup>1</sup> almost axiomatic.

Essential to normal cell metabolism and wound healing, these poorly-stored, readily-diffusible factors must be replenished—usually by *massive dosage*—if tissue rehabilitation<sup>2</sup> and return to health<sup>3</sup> are to be expedited. • Allbee with C 'Robins' provides this all-important "saturation dosage" in convenient capsule form. It incorporates the important B factors in 2 to 15 times daily requirements, plus 250 mg. of vitamin C—the *highest strength of ascorbic acid available today in a multi-vitamin capsule*. • Its prescription represents a sound contribution toward decisive recovery from disease, or toward pre- and post-operative nutritional support.<sup>4</sup>

**A. H. ROBINS CO., INC. • RICHMOND 20, VA.**  
*Ethical Pharmaceuticals of Merit since 1878*

**FORMULA:** Each Allbee with C capsule contains:

Thiamine hydrochloride (B <sub>1</sub> ).....	15 mg.
Riboflavin (B <sub>2</sub> ) .....	10 mg.
Nicotinamide .....	50 mg.
Calcium pantothenate .....	10 mg.
Ascorbic acid (C).....	250 mg.

**REFERENCES:** 1. Collier, F. A. and DeWesse, M. S.: Preoperative and Postoperative Care, J.A.M.A., 141:641, 1949. 2. Jolliffe, N. and Smith, J. J.: Med. Clin. North America, 27:567, 1943. 3. Kruse, H. D.: Proc. Conf. Convalescent Care, New York Acad. Med., 1940. 4. Spies, T. D.: Med. Clin. North America, 27:275, 1943.

**allbee® WITH C**



**An Important Advance**  
**IN THE PREVENTION AND TREATMENT OF**  
**IMPETIGO, EXCORIATED BUTTOCKS,**  
**MILIARIA, DIAPER RASH**

**—New Formula Johnson's Baby Lotion**

For more than 3 years, the Medical Research Laboratory of Johnson & Johnson in co-operation with several leading universities, has been engaged in fundamental research leading to the development of a physiologically acceptable preparation for use on infant's skin. These efforts have culminated in the release of the new Johnson's Baby Lotion following an accumulated experience of over 10,000 baby days.

The findings indicate that the new Johnson's Baby Lotion is a specific preventative and therapeutic agent for the five most common skin afflictions of infancy: impetigo contagiosa, miliaria rubra, intertrigo, excoriated buttocks, and diaper rash.

**Description and Pharmacologic Action**

Johnson's Baby Lotion consists of a nontoxic, nonirritating oil-in-water emulsion, which, when placed upon the skin, produces a discontinuous film having the ability to protect the skin from external irritative agents, but without interference with the transpiration of water vapor and other physiologic functions of the skin.

Johnson's Baby Lotion, by virtue of its bacteriostatic and bactericidal properties, produces a marked

and prolonged suppression of the resident bacterial flora of the skin, thus offering a substantial degree of protection against superficial infection.

**Clinical Evidence**

In 8 large hospitals, under the guidance of pediatricians and dermatologists, clinical investigations have been conducted on the new Johnson's Baby Lotion containing hexachlorophene\* in a concentration of 1% as an antiseptic agent. Herewith are pertinent excerpts from the reports. (Complete reports available on request.)

(\*Hexachlorophene has been adopted by the Council on Pharmacy and Chemistry of the American Medical Association as the generic designation of Dihydroxyhexachlorodiphenyl Methane.)

**In a Pennsylvania Hospital:** "Conclusive evidence has been obtained that the hexachlorophene lotion is less irritating than ammoniated mercury, commonly used in newborn nurseries, and is more effective in preventing the minor skin irritations and superficial infections common to the newborn."

In another Pennsylvania hospital: "In the height of an epidemic of impetigo the hexachlorophene lotion not only prevented babies from developing lesions, but on

those babies who were affected, the lesions were few, discrete, and disappeared quickly without any other therapy. The epidemic of impetigo, which had been continuing for four months, ceased within a period of a week to ten days after the lotion was used on all babies in the nursery.

"It was concluded that the lotion exhibited an antibacterial effect which was sufficient to modify remarkably the course of a virulent epidemic of impetigo contagiosa."

**In a New York State Hospital:** "The hexachlorophene lotion was found to be unusually satisfactory in the routine care of the skin of infants beyond the newborn period and to be prophylactically effective in minimizing the incidence of diaper rash and miliaria."

**In a Nebraska Hospital:** "We saw no evidence of irritation from Johnson's Baby Lotion either in the babies on whom the lotion was applied or among the nurses applying the Lotion. We did not see at any time during our work any sensitivity to Johnson's Baby Lotion and on some of our children the

Lotion has been applied at various times for a period of four months."

### Summary

Clinical evidence indicates that the new formula of Johnson's Baby Lotion, containing hexachlorophene, is outstandingly effective in the prevention and cure of the major skin afflictions of infancy: impetigo contagiosa, miliaria rubra, intertrigo, excoriated buttocks, and diaper rash. Free samples of Johnson's Baby Lotion are available for your examination and for distribution to patients.



## JOHNSON'S BABY LOTION

**FREE!**

Mail coupon  
for 12 distribution  
samples.

Johnson-Johnson

Johnson & Johnson  
Baby Products Division  
Dept. M 3, New Brunswick, N. J.

Please send me, free of charge, 12  
distribution samples of Johnson's Baby  
Lotion.

Name

Street

City  State

Offer limited to medical profession  
in U. S. A.

## SHORT REPORTS



### OPHTHALMOLOGY

#### Radiation Cataracts

Ocular defects have been noted in persons exposed to either atom bomb or laboratory radiation. The cyclotron-induced cataracts are probably the result of chronic exposure to neutron irradiation, believe Drs. P. H. Abelson of Carnegie Institution of Washington, D.C., and P. G. Kruger of University of Illinois, Urbana, since gamma-ray intensity is not high enough to cause cataracts. In a study of these defects in survivors of Hiroshima and Nagasaki, however, Dr. David G. Cogan of Harvard University, Boston, and associates of the University of California, San Francisco, were unable to determine the exact role played by either neutrons or gamma rays, both of which were present in the zone. Of the 10 scientists with ocular defects after exposure to laboratory radiation, only 3 suffered severe visual damage. The

others had insignificant to moderate changes with little or no loss of sight. Blood of most of the 10 had been examined periodically while working in high-voltage laboratories, but no blood changes were observed. The estimated exposure of the scientists was surprisingly low, from 10 to 135 n with a median of 50 n.

*Science 110:654-657, 1949.*

### HORMONES

#### Eosinophil Response to ACTH Therapy

Eosinophils completely disappear from the peripheral circulation of patients given intramuscular adrenocorticotrophic hormones. No eosinophils were found in the blood of 10 patients six to seventy-two hours after start of ACTH administration, although counts had averaged 472 cells before therapy. Dr. Theron G. Randolph and associates of Chicago report that eosinophils return to pre-treatment levels three to six days after cessation of therapy.


*Proc. Central Soc. Clin. Research 22:67-68, 1949.*

### STATISTICS

#### Surgery in the U.S.

General practitioners and nonsurgical specialists perform about four times as much of the surgery as the surgeons do. Of the 10,000,000 operations performed in the United States during 1949, the market research department of MODERN MEDICINE finds that general practitioners did about 45%, while specialists in surgery did less than half this amount. About one-fifth of all surgical operations are done outside the hospital—in the doctor's office, the patient's home, or elsewhere.

NOT A REVOLUTION  
but the *Evolution* of  
sound arthritis therapy



A ROERIG PREPARATION

**EACH CAPSULE CONTAINS:**

Vitamin D (Irradiated Ergosterol).....	50,000 U.S.P. Units
Vitamin A (Fish-Liver Oil).....	5,000 U.S.P. Units
Vitamin C (Ascorbic Acid).....	75 mg.
Vitamin B <sub>1</sub> (Thiamine Hydrochloride).....	3 mg.
Vitamin B <sub>2</sub> (Riboflavin).....	2 mg.
Vitamin B <sub>6</sub> (Pyridoxine Hydrochloride).....	0.3 mg.
Niacinamide.....	15 mg.
Calcium Pantothenate.....	1 mg.
D-alpha Tocopherol Acetate.....	2.4 mg.

(Equivalent by biological assay to 3.3 mg.  
International Standard Vitamin E)



Since the time of Hippocrates, revolutionary "cures" for arthritis have aroused the false hopes of countless sufferers. But the modern and generally accepted concept of systemic treatment of arthritis has evolved from years of painstaking investigation. During these years the Darthronol formula was gradually developed.

Each of the nine active constituents of Darthronol has been studied experimentally and clinically and is known to be essential for the optimal well-being and maximal functional efficiency of arthritic patients.

The return to gainful occupation of thousands of arthritics, who have taken Darthronol as part of a systemic rehabilitation regimen, is evidence of the efficacy of Darthronol in abolishing pain, diminishing soft tissue swelling and restoring useful function.

## DARTHRONOL

FOR THE ARTHRITIC

**J. B. ROERIG AND COMPANY**  
535 Lake Shore Drive, Chicago 11, Ill.

## SHORT REPORTS

### ONCOLOGY

#### Cancer Caused by Plastics

Cellophane and polyethylene implants may cause malignant growths within the body. Dr. B. S. Oppenheimer and associates of Columbia University, New York City, find that cellophane wrapped around the kidneys of white rats or embedded in the abdominal wall induces cancer in 35% of the animals. Such tumors, mostly fibrosarcomas, appear spontaneously in only about 1% of rats. Similar use of polyethylene produces sarcomas in more than 11% of animals. Cancers have not been observed after use of these plastic materials in human beings.

### AWARDS

#### Chemistry Honor

For research on the calcium chemistry of bone, Dr. Pauline Berry Mack of Pennsylvania State College will receive the 1950 Francis P. Garvan Medal honoring women in chemistry, announces the American Chemical Society.

### HOSPITALS

#### Hospital Appointments for Negro Physicians

The status of the Negro physician in New York City has improved within the past few years, according to an editorial in *New York Medicine*. In 1944, Dr. Conrad Berens, president of the Medical Society of the County of New York, took a firm stand for admission of Negro students to medical colleges, for acceptance of Negro physicians as members of hospital staffs, and against discrimination in admission to or certification by national medical or

surgical organizations. Since that time, advances have been made. The American College of Surgeons which had only 1 Negro member in 1944 now has more than 20. A negro has been elected to the House of Delegates of the American Medical Association, and 170 Negro physicians hold 258 different appointments in 32 New York hospitals. More than 90% of the city's Negro physicians under forty years of age now have hospital staff appointments.

*New York Med. 5:13-14, 1949.*

### NUTRITION

#### B<sub>12</sub> and Child Growth

Daily oral doses of vitamin B<sub>12</sub> are apparently effective in fostering growth of children. Although recognized as a growth-promoting agent for bacteria and animals, the vitamin has been used in man chiefly for treatment of pernicious anemia and sprue. The nutritional value of B<sub>12</sub> was studied by Dr. Norman C. Wetzel and associates at the Children's Fresh Air Camp and Hospital, Cleveland. A group of 11 children under regular care for malnutrition and slow growth were given 10 µg of crystalline vitamin B<sub>12</sub> orally in addition to whatever treatment they were having at the time. This single change in routine brought dramatic results for 5 of the children. Not only were their growth rates accelerated, but physical vigor, mental attitude, and appetite improved. The most striking results were obtained for a boy with allergic bronchitis. Besides growth improvement his asthmatic symptoms disappeared in the first week of B<sub>12</sub> therapy.

*Science 110:651-653, 1949.*





## for emotional equilibrium in the menopause

BENZEBAR\* not only frequently alleviates the depression you see in menopausal patients, but also the nervousness.

'BENZEBAR' is a logical combination of Benzedrine\* Sulfate (racemic amphetamine sulfate, S.K.F.) and phenobarbital. Thus, it provides the unique improvement of mood characteristic of 'Benzedrine' Sulfate and the mild sedation of phenobarbital. These two established agents work together to stabilize the patient's emotions and to restore her zest for life and living. *Smith, Kline & French Laboratories, Philadelphia*

# Benzebar

for the depressed  
and nervous patient

\*'Benzedrine' and 'Benzobar' T.M. Reg. U.S. Pat. Off.





# NOW AVAILABLE..

## BALANCED HORMONE THERAPY

**Estrogen plus  
Androgen in a  
Single Injection  
For Dual Pituitary  
Inhibition**

### DI-STEROID

TRADEMARK

BRAND OF ESTROGENIC  
SUBSTANCE AND TESTOSTERONE

*Kremers • Urban*

IN **DIPHASOL**

*I*n several conditions in the female in which estrogens and androgens are useful separately, the mechanism of action appears to be inhibition of the pituitary; **DI-STEROID\*** supplies the synergistic pituitary-inhibiting action of both hormones, thus permitting smaller doses of each and reducing the risk of side-effects.

#### FEATURES

● Avoids masculinizing effects ● Obviates disagreeable hyper-estrogenic effects (breast and pelvis tenderness and withdrawal bleeding) ● Valuable in conditions requiring treatment over relatively long periods ● In **DIPHASOL** solution—provides the steroids for both immediate and prolonged effects.

**INDICATIONS:** Menopausal syndrome; suppression of lactation and painful breast engorgement post partum; mastopathies; menometrorrhagia; and dysmenorrhea.

**SUPPLIED:** 10-cc. multiple-dose vials; 1-cc. ampuls, boxes of 6 and 25.

*Kremers • Urban Company*

PHARMACEUTICAL CHEMISTS SINCE 1894



# UNIQUE INJECTION SOLVENT...

**Permitting Immediate  
and Prolonged  
Hormonal Effect**

## DIPHASOL

PATENT PENDING

**DIPHASIC VEHICLE**

*Kremers • Urban*

**D**IPHASOL<sup>1</sup> solutions of hormones are clear, non-oily . . . no crystals to clog the needle.

DIPHASOL solutions replace conventional suspensions of hormones. Microcrystals are deposited in the muscle fibres only after dilution in situ by the tissue fluids.

**HOW IT WORKS:** Upon injection of a DIPHASOL solution of steroids, part of the solution passes immediately into the lymphatic circulation for rapid hormone effect; the remainder, when diluted by the tissue fluids, deposits microcrystals which are absorbed more slowly, for prolonged effect.

Kremers-Urban specialties available in DIPHASOL solution include ESTRUGENONE,\* PROSTRUGEN,\* DI-STEROID,\* DI-STERONE\* (testosterone and estrogens, testosterone predominating), DI-ERONE\* (testosterone and progesterone), testosterone, and progesterone.

DIPHASOL contains propylene glycol, ethyl alcohol, isotonic solution of sodium chloride, and benzyl alcohol.

1. Inject with a 26-gauge needle.
2. Gives prompt and prolonged effect.
3. Clean syringes with water.

\*Trademark of Kremers-Urban Co.

**BOX 2038 • MILWAUKEE 1, WISCONSIN**

## SHORT REPORTS

### METABOLISM

#### Parenteral Nutrition

Intravenous administration of a 15% emulsion of fat may prevent weight loss and help maintain a positive nitrogen and potassium balance when illness impedes adequate nutrition. The emulsion, consisting of 15% coconut oil, 4.3% dextrose, and a combination of 0.5% soybean phosphatides and 1% polyglycerol esters, is administered at rates used for glucose and saline solutions and supplies 1,600 calories per liter. Daily amounts of 3 and 6 gm. of fat per kilogram of body weight administered to adults and infants, respectively, maintain weight and positive nitrogen and potassium balance. In 3 patients, Dr. Sherwood W. Gorens and associates of Harvard University, Boston, found that amounts ranging from 0.8 to 4 gm. of fat per kilogram of body weight caused a rise of 1 to 3 degrees in temperature. Other patients received the same amounts or more, however, with no such effect. Pyrogenic materials in the fat rather than the fat itself were probably responsible. The emulsions do not appear to cause pathologic change.

*J. Lab. & Clin. Med.* 34:1627-1633, 1949.

### ENDOCRINOLOGY

#### Cortisone in Male Urine

The compound 17-hydroxy 11-dehydro corticosterone can be isolated from the urine of normal males. Dr. John J. Schneider of Jefferson Medical College, Philadelphia, finds that the material, cortisone, can be procured from a chloroform extract of pooled normal human urine obtained before hydrolysis.

*Science* 111:61, 1950.



"It may look like 'A' to you but I calls 'em as I sees 'em!"

### VITAMINS

#### Lipotropic Effect of B<sub>12</sub>

When given to rats in conjunction with a high-fat diet, vitamin B<sub>12</sub> concentrate has a lipotropic effect similar to that of liver extract. After twenty-nine days, Drs. Victor A. Drill and Harry M. McCormick of Wayne University, Detroit, found that the average fat content of livers of animals given 0.2 µg of vitamin B<sub>12</sub> three times a week was only about 16%. When the vitamin was withheld, the fat content was nearly twice as much. The treated animals also gained more weight than untreated rats on the same diet. When 1 µg was given, the amount of liver fat was approximately the same as in rats fed a normal diet. The exact nature of the lipotropic effect of vitamin B<sub>12</sub> is not known. The concentrate contains insufficient choline for that element to be responsible.

*Proc. Soc. Exper. Biol. & Med.* 72:388-390, 1949.

# PROTAMIDE

FOR THE LIGHTNING  
PAINS AND ATROXIA OF  
**TABES  
DORSALIS**



**A DRAMATIC THERAPY  
CLINICALLY PROVED FOR  
THE RELIEF OF PAIN  
AND LESIONS OF  
HERPES ZOSTER**



AVAILABLE IN  
10 AMPUL 1.3 cc. SIZE.  
PHYSICIAN'S PRICE  
\$15.00

**SEND FOR LITERATURE:** Please write for literature and reprint... Protamide is available from your local supplier, or direct from us. Protamide is stable at room temperatures. Keep a package on hand... Gain experience with this valuable contribution to therapy.

*\*Extensive Clinical Data on Request.*

**DESCRIPTION:** An aqueous, colloidal solution of a proteolytic enzyme, processed by the original method discovered and developed by Fuller. Each lot is clinically assayed for efficacy.

**INDICATIONS:** Critical evaluation by competent investigators has firmly established Protamide as the therapy of choice for herpes zoster and tabes dorsalis.

**ADMINISTRATION:** Freedom from side effects and comparative absence of pain on intramuscular injection—adds to the acceptance of Protamide.

**DOSAGE AND RESULTS:** The pain of herpes zoster is often relieved after the first injection. However, an ampul daily for 3 to 4 days is recommended. Pain is controlled and lesions heal rapidly in the great majority of cases. Of 45 patients with tabes dorsalis receiving 12 ampuls or more of Protamide, 44 received relief from lightning pains.\*

**SHERMAN LABORATORIES**  
G. H. Sherman, M. D., Founder  
**BIOLOGICALS • PHARMACEUTICALS**  
DETROIT 15, MICHIGAN

REGISTERED U. S. TRADE MARK

## SHORT REPORTS

### CIRRHOSIS

#### Serum Lipids and Xanthomas

Occurrence of skin xanthomas with primary biliary cirrhosis seems to be directly related to the degree of elevation of the total serum lipids. When the elevation is above 2,000 mg. per cent for a prolonged period of time, Drs. Edward H. Ahrens, Jr., and Henry G. Kunkel of the Rockefeller Institute for Medical Research, New York City, find that skin xanthomas usually develop. If the biliary obstruction is relieved, the lipids fall and the xanthomas resolve. If the level does not exceed 1,800 mg. per cent, the patient will probably remain in the pre-xanthomatous phase of the disease. The lipid pattern of primary biliary cirrhosis comprises notable elevation of the phospholipids and lesser but significant increases of free cholesterol and neutral fat, with clear nonlipemic serum.

*J. Clin. Investigation* 28:1565-1574, 1949.

### SURGERY

#### Portal Occlusion

Irritative cellophane may be useful in preparing the portal vein for ligation and in experimental study of decreased blood flow. The vein is encased in a double cuff of Polythene and the diameter is then tightened one-third to one-half by a narrow tantalum band. By this method, Drs. Peter W. Stone and Ralph A. Murphy, Jr., of Emory University, Atlanta, Ga., produced partial occlusion in dogs with development of adequate collateral circulation in five days and complete obliteration of the lumen in forty to sixty days.

*Proc. Soc. Exper. Biol. & Med.* 72:255-259, 1949.

### PREVENTIVE MEDICINE

#### Fluorine-Goiter Relationship

The addition of fluorine to drinking water for the prevention of tooth decay may increase the incidence of goiter. Dr. D. G. Steyn of the University of Pretoria has found such a trend in several regions of South Africa where fluorine is being added. The material, which has chemical properties similar to those of iodine, is attracted to the thyroid and acts to deprive the gland of iodine. This action is intensified by water with a high-calcium content. Goiters may be prevented in persons drinking such water by a greater intake of iodine.

### DIAGNOSIS

#### Neurogenic Appendicitis

Abdominal pain in the right lower quadrant, with nausea, vomiting, and possibly fever, may be due to a tumor-like growth of nerve fibers and ganglion cells in the appendix. In 536 vermiform processes examined by Gustav Lassmann, M.D., of Vienna, 91 cases of appendicular neuromas were found. Pathologic findings were limited to the nerve endings. Secondary inflammation was found in only 5 of the 91 cases. Postoperative observation of 16 patients revealed that 6 were not relieved by surgery. The appendix should be removed early, while the process is still localized.

*Mikroskopie* 4, 9/10:277-290, 1949.

### WHO


#### World Health

Peru recently joined the World Health Organization, becoming its sixty-seventh member.

# Stop-Gap for Headache Pain **Anacin**



While the cause of the headache is being determined, the patient can enjoy relief from pain with the aid of Anacin. This dependable APC formula not only gives fast relief, but long-lasting relief as well, exceeding that of plain aspirin. When your patients need fast, prolonged relief of headache, neuritis or neuralgia pain, why not recommend Anacin tablets? Available at all drug stores and hospital pharmacies.

**ANACIN**

WHITEHALL PHARMACAL COMPANY • 22 East 40th Street, New York 16, N. Y.



## SHORT REPORTS

### HORMONES

#### Anaphylactoid Reaction

Both cortisone and ACTH have a strongly inhibiting effect on the anaphylactoid edema caused by the par-enteral administration of egg white to rats. Desoxycorticosterone and lyophilized anterior pituitary tend to aggravate the reaction. Dr. Hans Selye of the Université de Montréal, Canada, has found that gluco-corticoids such as cortisone inhibit many of the actions of mineralo-corticoids such as DCA. This observation suggests that adrenalectomy sensitizes the patient to the toxic reactions of DCA by predisposing to an unfavorable gluco-corticoid and mineralo-corticoid balance. Lesions particularly amenable to ACTH or cortisone usually react favorably to non-specific therapy, fever therapy, and shock therapy. Since these measures stimulate the production of gluco-corticoids, many of the so-called collagen diseases may depend on a derangement of the pituitary-adrenocortical response to stress.

*Canad. M. A. J.* 61:553-556, 1949.

### HORMONES

#### Rheumatoid Arthritis Therapy

Combined injections of desoxycorticosterone acetate and ascorbic acid have been used in the treatment of rheumatoid arthritis with satisfactory results. Fifteen to thirty minutes after injections, pain almost completely disappeared and mobility improved as much as anatomic joint changes and muscular atrophy would allow for all of 9 patients with slight or severe involvement of two weeks' to fifteen years' duration. Effects last two to six hours, sometimes a day,

and are apparently more protracted after each subsequent injection. The pain-relieving effect appears to be confined to the joints. No exact dosages have been worked out, but Drs. E. Lewin and E. Wassén of Sahlgren's Hospital, Gothenburg, Sweden, observe that increasing the dose to more than 5 mg. of DCA with or without 1 gm. of ascorbic acid does not seem to enhance effects, and results are almost equally good with half these amounts. DCA is usually injected intramuscularly and ascorbic acid intravenously two to five minutes later. A delay of as long as two hours between the injections renders the treatment ineffective. No toxic side effects have been observed. Observation time has been two or three weeks.

*Lancet* 257:993, 1949.

### VIRUS DISEASE

#### Antibiotics for Psittacosis

Life of a chick embryo infected with psittacosis virus can be prolonged by either aureomycin or Chloromycetin, but the latter drug is somewhat less effective. Using equal doses of each antibiotic, Drs. E. Buist Wells and Maxwell Finland of Harvard University, Boston, find that embryos protected with aureomycin live two to three days longer than those given Chloromycetin. Prolongation of life appears to be directly related to dosage for each drug. On a weight basis, aureomycin is more than 3 times as effective as Chloromycetin against psittacosis and more than 5 times as effective on a molecular basis.

*Proc. Soc. Exper. Biol. & Med.* 72:365-368, 1949.

# A.C.M.I. Portable ELECTRO-SURGICAL UNIT

No. C-350

Within the compact case of this newly designed, readily portable, inexpensive unit, A.C.M.I. engineers have packed a remarkably efficient and ruggedly dependable high frequency unit, amply powered for such electro-surgical procedures as cutting, coagulation, desiccation and fulguration.

The unit provides a vacuum tube cutting outlet, and a spark gap circuit for coagulation, desiccation and fulguration. A selector switch, centrally located, permits quick and unerring selection of tube cutting, coagulating or blended current, as desired. Tube cutting intensity is readily controlled by the right-hand knob, while the spark gap control (for coagulation, desiccation or fulguration) is conveniently located on the left of the panel. Electrodes, cables and other accessories, are carried in the cover of the unit, which is mounted in a durably handsome leatherette case, weighing complete only 32 pounds.

Write for literature

**American Cystoscope Makers, Inc.**

1341 LAFAYETTE AVE.  
FREDERICK J. WALLACE



NEW YORK 39, N. Y.  
PRESIDENT



Complete unit  
supplied in  
light weight  
leatherette case

Accessories carried  
handily in cover



## The Accident-prone Automobile Driver

W. A. TILLMANN, M.D., AND G. E. HOBBS, M.D.\*

*University of Western Ontario, London, Ont.*

**A** RELATIVELY small group of drivers seems to be responsible for a large number of automobile accidents.\*

In a study of high- and low-accident groups among the general driving population and drivers of commercial vehicles, W. A. Tillmann, M.D., and G. E. Hobbs, M.D., find a striking degree of similarity in personalities and backgrounds of the accident-prone drivers. A man's conduct behind the wheel of a car appears to simulate his behavior in other situations.

In general, the high-accident driver has a record of instability and resistance to authority. His parents are probably incompatible and may have been divorced while he was a child.


One or both parents were usually excessively strict. Childhood instability was expressed either in aggressiveness, bullying, and juvenile delinquency or in extreme introversion and many hidden fears. In school he was probably an average student,

but his record was marred by truancy and breeches of discipline.

After the accident-prone driver went to work, he held many different jobs in a short time and was often discharged. If a taxi driver, he probably prefers such work to a more settled routine. As a rule, gambling, dancing, and sports are his principal recreations.

He usually has many acquaintances but few close friends. Sexual promiscuity is common and usually freely admitted and, if married, he evinces little interest in his family. His conversation is frequently interesting but usually compounded of cursing, ob-

*(Continued on page 104)*



\* The accident-prone automobile driver. *Am. J. Psychiat.* 106:321-331, 1949.

Now pre-calibrated for


**40** common clinical tests  
including Thymol Turbidity,  
Red Blood Cell Count

The ROUY

**Photrometer**

by *Leitz*

To make the Photrometer even more helpful to busy doctors, Leitz now pre-calibrates for Thymol Turbidity and Red Blood Cell Count procedures. Since pre-calibration eliminates calculation and the need for preparing standards, the Photrometer simplifies and speeds up 40 of the most commonly required clinical tests.



Unique improvements make the Photrometer the most highly selective and accurate photoelectric colorimeter on the market...accurate to within 0.1%. And it's so amazingly simple to operate! There are only two moving parts—a single control for all adjustments. An individual handbook accompanies each Photrometer and contains outlines of standard methods which any competent worker can easily follow. With each method there is a calibration table which eliminates any need for calculation.

Here is an invaluable lab assistant that will soon pay for itself in time and money saved. See for yourself what a timesaving aid it can be in a busy practice.

**Pre-calibrated for 40 tests—\$257.80**

**Pre-calibrated for 22 tests—\$182.80**

*For particulars, see your Leitz franchised dealer*

**E. LEITZ, Inc.**

**304 Hudson Street, New York 13, N. Y.**

### TRADE-INS

Old model Leitz Colorimeters will be accepted by your Leitz franchised dealer when you buy the new Rouy Photrometer.

LEITZ SCIENTIFIC INSTRUMENTS • MICROSCOPES • LEICA CAMERAS AND ACCESSORIES

in  
rheumatic  
affections...



*how can Pabalate provide  
better salicylate therapy  
than pure salicylate itself?*

**THE SUCCESS** of salicylate therapy in rheumatic affections has been shown by authoritative reports<sup>3,4</sup> to depend largely on the maintenance of really adequate blood levels . . . frequently a difficult achievement under usual salicylate administration. Pabalate supplies not only salicylate, but also a "booster" in the form of the antirheumatic para-aminobenzoic acid,<sup>7</sup> which acts to increase blood levels of salicylate.<sup>1,2,4,5</sup> In turn, the salicylate increases the blood concentration of the para-aminobenzoic acid.<sup>2</sup> Enteric coating helps Pabalate prevent gastric irritation, insures optimal toleration. Successful clinical results, contingent on adequate blood levels, can thus be achieved better, more dependably, with Pabalate . . . the "new word for salicylate" in therapy of rheumatic affections.

**A. H. ROBINS COMPANY, INC. • RICHMOND 20, VA.**  
Ethical Pharmaceuticals of Merit since 1878

higher salicylate blood levels for better antirheumatic therapy

**INDICATIONS:**

**Pabalate Tablets**—for adult patients with rheumatoid arthritis, acute rheumatic fever, fibrositis, gout and osteo-arthritis. **Liquid Pabalate**—for treatment of acute rheumatic fever or other rheumatic diseases in children and as a replacement for tablet salicylate medication; or for adults who prefer a liquid dosage form.

**DOSAGE:**

Average adult dose: two tablets or teaspoonfuls, three or four times daily. Dosage should be adjusted upward if necessary. For children, dosage is proportional to age and severity of condition.

**FORMULA:**

Each enteric-coated tablet or each teaspoonful contains Sodium Salicylate, U.S.P. (5 grs.) 0.3 Gm.; Para-aminobenzoic Acid (as the sodium salt) (5 grs.) 0.3 Gm.


**SUPPLIED:**

**Pabalate Tablets** in bottles of 100 and 500. **Liquid Pabalate** in bottles of 1 pint.

**REFERENCES:**

1. Bellisle, M.: *Union Med. Canada*, 77:392, 1948
2. Dry, T. J. et al.: *Proc. Staff Meetings Mayo Clinic*, 21:497, 1946
3. Editorial: *J.A.M.A.*, 138:367, 1948
4. Muratore, F. and Pugignone, T.: *Bull. Soc. Ital. Biol. Sper.*, 24:269, 1948
5. Parker, W. A.: *Quart. J. Med.*, 17:229, 1948
6. Reid, J.: *Quart. J. Med.*, 17:129, 1948
7. Rosenblum, H. and Fraser, L. E.: *Proc. Soc. Exper. Biol. and Med.*, 65:178, 1947

**Para-aminobenzoic acid increases blood levels of concurrently administered salicylate.\***



For treatment of rheumatic affections

Rx *Pabalate*

Robins



## MEDICAL NEWS

scenity, and obvious attempts to impress his listeners. Personal dress tends to be eccentric.

His health record shows the common childhood diseases and a number of personal injuries from accidents, but few functional complaints.


On the road his principal concern is to bluff the other driver. He uses the horn a great deal and is particularly angered at finding his own faults in other motorists. Interest in the gadgets and mechanical aspects of the car, but no concern for mechanical limitations is another mark of the accident-prone driver.

Finally, his outlook on life is likely to be fatalistic and materialistic. He lives from day to day with no thought for the future.

The personal history of the driver with few accidents is quite different. His parents were usually congenial and were firm but understanding disciplinarians. As a child he may have been slightly unstable, but not aggressive. His academic standing was average, but disciplinary infractions were few.

This man usually has worked at one job for a long period of time and, if a taxi driver, he would probably prefer employment that offers more security. His interests and recreational activities tend to be conservative, including little drinking and almost no gambling. He has a large circle of friends.

Sexual promiscuity is uncommon and any dereliction is accompanied



### For governed maintenance...

Digitaline Nativelle maintains the maximum efficiency obtainable. *Positive maintenance*—because absorption is complete and the uniform rate of dissipation provides full digitalis effect between doses. All, with virtual freedom from side reactions.

## DIGITALINE NATIVELLE

Chief active principle\* of digitalis purpurea [digitoxin]

\*Not an adventurous mixture of glycosides.

**MAINTENANCE:** 0.1 or 0.2 mg. daily depending on patients' response. **CHANGE-OVER:** 0.1 or 0.2 mg. Digitaline Nativelle replaces 0.1 or 0.2 gm. whole leaf. **RAPID DIGITALIZATION:** 0.6 mg. initially, followed by 0.2 or 0.4 mg. every 3 hours until patient is digitalized.

Send for brochure "Modern Digitalis Therapy" Varick Pharmacal Co. Inc. (Div. E. Fougere & Co. Inc.), 75 Varick St., New York

*simplicity, itself*



*to prescribe* **SIMILAC**

simply add one measure of Similac to  
two ounces of water to yield two ounces  
of normal formula of 20 cal/oz

*simplicity, itself*



*to prepare* **SIMILAC**

simply instruct mother to float the  
prescribed quantity of Similac  
on previously boiled water and stir

*simplicity, itself*



*to digest* **SIMILAC**

*for* { the proteins have been so modified  
the fats so altered  
the minerals so adjusted

*that there is no closer equivalent  
to human breast milk than*

**SIMILAC**

for term and premature infants throughout the first  
year of life whenever breast feeding must be  
supplemented or replaced. Similac has the same zero  
curd tension as human breast milk.

SIMILAC DIVISION



M & R DIETETIC LABORATORIES, Columbus 16, Ohio

## MEDICAL NEWS

by a feeling of guilt. This driver is probably quiet and reserved; he may appear unsociable. His health is probably not as good as that of the accident repeater.

His driving habits are courteous, and he is on the alert for other drivers who may make mistakes. Usually he has no interest in mechanics, is concerned with the welfare of others, and worries a great deal.


In most cases the accident tendency seems to be a fixed trait, but may be reversed by a change in the emotional pattern of the individual. A taxi driver with the background and the personal characteristics of a typical accident-prone individual drove 50,000 miles with 2 serious and 6 minor accidents between the ages

of sixteen and twenty-six. After marrying a girl with strong religious inclinations, his behavior changed, and he drove about 550,000 miles in the next twelve years with only 1 minor accident.

Studies have not shown a significant relation between driving skills and accident rate. Safe driving depends more upon judgment and caution than upon mechanical skills. In accident-proneness, the defect is always above, and never below the basal ganglia.

If a personal history were added to driving tests, the accident-prone driver could be discovered even before his driving record unmasks him, and only safe drivers would be licensed.

**an effective sedative and prompt-acting hypnotic**



Clinical experience has demonstrated the efficacy of NEURONIDIA in securing sedation and hypnosis.

Neuronidia offers barbital that has been pleasantly masked in a palatable elixir to provide complete flexibility of dosage.

In bottles of 8 fl. oz.

**Schieffelin & Co.**

Pharmaceutical and  
Research Laboratories  
20 Cooper Square,  
New York 3, New York

**neuronidia**



*For the weary  
old woman  
of 21*



*Three  
Iberol Tablets supply:*

FERROUS SULFATE, U.S.P. 1.05 Gm.  
(representing 210 mg. elemental  
iron, the active ingredient for the  
increase of hemoglobin in the treat-  
ment of iron-deficiency anemia.)  
plus these nutritional constituents:

Thiamine Hydrochloride ..... 6 mg. (6 x MDR\*)  
Riboflavin ..... 6 mg. (3 x MDR\*)  
Nicotinamide... 30 mg. (2 x RDA†)  
Ascorbic Acid 150 mg. (5 x MDR\*)  
Pyridoxine Hydrochloride... 3 mg.  
Pantothenic Acid ..... 6 mg.  
(as Calcium Pantothenate)  
Folic Acid ..... 5.1 mg.  
Liver Concentrate, N.F. ... 1.5 Gm.

\*Minimum Daily Requirement  
†Recommended Daily Dietary Allowance



**S**HE comes to you  
with marked pallor, complaining of  
chronic fatigue, no appetite. Her blood  
picture will probably reveal iron deficiency  
anemia, and judging by her lack of appetite and  
general condition you may suspect a concurrent vitamin  
deficiency. • In such cases, isn't it more convenient and  
economical to correct both deficiencies with a single  
preparation? Just three **IBEROL** tablets, one with each  
meal, supply 210 mg. elemental iron plus generous  
amounts of ascorbic acid, vitamin B complex and  
liver concentrate. Yet, in spite of such potency,  
**IBEROL** tablets are compact and easy to swallow.

The triple coating makes this possible—Abbott  
employs the ferrous sulfate itself as one of the  
coatings, thus reducing bulk and separating  
the iron from vitamins with  
which it is not compatible.

**Abbott**

***Iberol*** <sup>®</sup> **TABLETS**

(IRON-VITAMIN B COMPLEX-LIVER CONCENTRATE, ABBOTT)

*Without Folic Acid* **MACRON\* TABLETS**

have the same potent formula as Iberol,  
but with the folic acid omitted.

•TRADE MARK

# Washington Letter

## An Expensive Paradox in the Federal Medical Program

The administration's budget for the next fiscal year, now on its way through Congress, brings into clear focus the most costly paradox in the federal medical program.

Medical facilities are being constructed to care for veterans who have non-service connected disabilities, while at the same time administration leaders join nonpartisan crit-

ics in condemning expansion of this medical care system.

In this election year, the administration apparently is not demanding change. In his budget message, the President merely skirted this issue. Marshaling arguments against costly benefits, Mr. Truman said:

I again urge that, in considering new or additional aids for veterans without service disabilities, Congress judge their necessity not merely from the standpoint of military service, but also on the basis of benefits under the general social security, health and education programs available to all people, including veterans. . . . We should provide only for the special and unique needs of veterans arising directly from military service.

However, but a few hundred words farther along in his message, Mr. Truman revealed that Veterans Administration is preparing to handle more, not fewer, cases of non-service connected injuries or illnesses. On this point, the President says:

Construction of hospitals to provide 37,000 new beds and addition-



"Now this may hurt just a little bit, Doc."



**after tonsillectomy...**

the chewing of Aspergum,  
before mealtime, effectively  
relieves pain, lessens muscle  
spasm, contributes to  
patient comfort.

DILLARD'S

aspergum

**SALIVARY ANALGESIA**

Contains 3 1/2 grains of  
aspirin in a pleasantly  
flavored chewing gum base—  
particularly suitable for  
administering aspirin to  
children and to patients who  
have difficulty swallowing tablets.  
*Ethically promoted.*

Aspergum is available  
at all prescription  
pharmacies in  
bottles of 36 and 250,  
packages of 16.

**WHITE LABORATORIES, INC.,**  
**Pharmaceutical Manufacturers,**  
**Newark 7, N. J.**



## WASHINGTON LETTER

al domiciliary facilities, costing \$872,000,000, is now about one-third completed. By June 1951 it is estimated that three-fourths of the work will have been done. When this program is finished, there will be sufficient beds to provide adequately for foreseeable needs for all service-connected cases and a more liberal allowance of beds than at present for non-service connected cases.

There is no suggestion that Congress reexamine the practice of allowing VA to hospitalize veterans for troubles that have no connection with military service.

Immediately following the above quotation, the budget message sets out in round millions of dollars what this service is costing. It states:

Current expenses for hospital and medical care are estimated at 590 million dollars in the fiscal year 1951 (starting July 1, 1950). About four-fifths of these expenditures are for the in-patient care program, and in this program two-thirds of the cases currently are non-service connected.

Incidentally, there is an additional cost from this service which VA budget does not reflect. Thousands of veterans with non-service connected disabilities are hospitalized in Army and Navy hospitals. Much of the cost of caring for them appears on

the defense budget, not the VA budget. Just what this costs is difficult to estimate, certainly many millions of dollars each year. The total is high enough to cause serious concern among some defense department medical leaders who recently induced Defense Secretary Louis Johnson to issue an order to military hospitals reminding the staff that the primary task was to care for military personnel. However, it is doubtful that this order will affect the present practice.

### President and Health Insurance

In none of his communications to Congress so far has Mr. Truman demanded passage this session of his national health insurance program. No one should be misled by the absence of such a demand. Inquiries at the White House clear up any doubt. Mr. Truman is still convinced that nothing but comprehensive, payroll deduction health insurance will provide the nation with adequate medical care. Ultimately, he thinks, Congress and the people will see it this way and legislation along the lines of the Wagner-Murray-Dingell bill will become law.

The impression from the White



# on the **COURSE** to bounding health...

## with Citrus Fruits and Juices

The rich variety of nutrient factors\* in citrus fruits and juices, and their high content of vitamin C and natural fruit sugars,<sup>1</sup> are medically noteworthy — for several reasons: they contribute helpfully to improvement of appetite<sup>2</sup> and digestion,<sup>3</sup> to greater bodily energy<sup>4</sup> and stamina,<sup>5</sup> and to resistance to disease.<sup>6</sup> During pregnancy or lactation, before or after surgery, and for general nutritional support from infancy to old age, the refreshing goodness of low-cost, readily-available Florida-grown fruits and their juices proves universally appealing ... whether in fresh, canned, frozen or concentrated form.

**FLORIDA CITRUS COMMISSION  
LAKELAND, FLORIDA**

\*Among the richest known sources of vitamin C are the citrus fruits. They also contain vitamins A, B, and P, and readily assimilable natural fruit sugars — together with other factors such as iron, calcium, citrates and citric acid.

*References:* 1. Gordon, E. S.: *Nutritional and Vitamin Therapy in General Practice*, Year Book Pub., 2nd ed., 1947. 2. Manchester, T. C.: *Food Research*, 7:284, 1942. 3. McLeskey, J. S.: *Nutrition and Diet*, Saunders, 4th ed., 1944. 4. Howe, H. S.: *How's Your Foundation of Nutrition*, rev. by MacLeod and Taylor, Macmillan, 2nd ed., 1944. 5. Sherman, H. C.: *Chemistry of Food and Nutrition*, Macmillan, 7th ed., 1948.



# **FLORIDA**

**Oranges • Grapefruit  
Tangerines**

## WASHINGTON LETTER

House is that Mr. Truman is not dismayed by lack of interest in his health bill by so many Democrats in Congress. He knows this means delay, but he's convinced it does not mean defeat.

### Left Over from Last Session

A number of important health bills were started through the last session of Congress but did not become law. Unless this session becomes tied up in long controversies over taxes, civil rights bills, or foreign aid, several of these health bills have a good chance of passing.

► The school health program, calling for \$25,000,000, is almost certain to pass; it nearly became law last session.

► Another bill that has good prospects is one that would increase grants for local public health assistance from \$14,000,000 to \$23,000,000.

► The bill to help finance medical and related schools also is in high favor in Congress; most of the controversial issues were compromised before end of the last session.

► Creation of a National Science Foundation in the next few months is almost assured; unusual legislative complications were the only reason it was not passed last session.

► If the administration's budget is followed, appropriations for education and general research will be more than trebled, rising from \$125,000,000 the current fiscal year to \$434,000,000.

## Krestex

### PLASTICALLY PERFECT PROTECTION!

Professionally Designed for Professional Use!

• Krestex professional garments are modern miracles—affording a perfect and practical protection never before possible. Krestex is impervious to staining—acid and chemical resistant—waterproof! Wiped with a damp cloth, it comes clean in a jiffy (eliminates laundry expense). Unequaled for quality and amazing durability. Unmatched for dignified professional appearance.

### Complete Line of KRESTEX Products for All Your Needs . . .

• Surgical Aprons • Nurses' Aprons  
• Laboratory Aprons • Equipment Covers  
• Sheets • Pillow Cases • Mattress Covers, etc.

### INSIST ON THE BEST—KRESTEX!

If Your Dealer Cannot Supply,  
Write Us Direct. Send for  
Brochure of  
Complete Line

*Petra*




### DOCTOR'S BOOTEES

Protect Trousers and Shoes from  
Plaster When Applying Casts  
Specially designed for wear  
during orthopedic procedures—  
also other uses where shoes and  
trousers are exposed to stains.  
Generously sized to slip over  
shoes easily.

(Prices Slightly Higher West of the Rockies)

**350**  
PAIR

MANUFACTURING COMPANY • Dept. MM • 1335 N. WELLS ST. CHICAGO 10, ILLINOIS



**\*PLUMS—BUT WHAT PLUMS!** Sugar-sweet purple beauties, strained to velvety purée. Then just a touch of Tapioca Flour (not over 5%) is added for the extra-smooth texture you and we know is a must with tiny infants. *New, too! Gerber's Junior Plums.* Finely chopped for tots with a few teeth.

**OTHER RECENT ADDITIONS** to the ever-increasing variety of delicious, nutritious Gerber's: Sweet Potatoes and True-Flavor Meats. We're working on others. For constantly helping you encourage good eating habits in infancy is our only business!

**SAMPLES** of Plums with Tapioca and Gerber's 3 Cereals. Write to Dept. 213-0, Fremont, Mich.



**Gerber's**  
BABY FOODS  
Fremont, Mich.



**Babies are our business...our only business!**

## WASHINGTON LETTER

### Rehabilitation Dividend

Statistics of the Office of Vocational Rehabilitation, covering a five-year period, show that as a nation we are not wasting money on the \$76,000,000 rehabilitation appropriation. The program has resulted in the rehabilitation of almost a quarter of a million men and women to the extent that they have become able to support themselves and in some cases their families.

It is calculated that these formerly handicapped people have added just slightly less than one billion dollars to the nation's income over the five years. They have paid \$70,000,000 in federal income taxes alone.

Here is one example of how federal-state rehabilitation works:

In 1948 the average cost of rehabilitating one person was \$460, which is paid only once. Yet it

costs \$492 to maintain a blind person in dependency for just one year. The report points out that more than a million and a half persons still are so seriously disabled as to require rehabilitation service if they are to become self-supporting. Currently, federal and state funds are adequate to care for only 1 out of 5 of the handicapped.


### Good Neighbor, Dr. Hyde

Dr. H. van Zile Hyde, who has contributed ability and energy to the World Health Organization, is the new health and sanitation director for the Institute of American Affairs. The Institute was created to carry on cooperative programs among fourteen of the Latin American republics. In another capacity, it is the regional organization through which WHO will function in these areas. . . . Public Health Service's Communicable Disease Center at Atlanta gave emergency aid to seventeen states where epidemics or disasters occurred during the last year. It assisted five states that had poliomyelitis epidemics and five others in which encephalitis broke out. The other emergencies were caused by floods. . . . The one-thousandth hospital has been approved for construction under the Hospital Act. This act was liberalized in the last session of Congress and offers greater opportunity to small towns and low-income communities. . . . Public Health Service will continue its influenza information center. This office, set up a year ago, is western hemisphere headquarters for a world-wide campaign for research into and control of this disease.



*"Don't be alarmed. I've just had a little trouble with the plumbing."*

# What do you look for in an antiarthritic?



## EFFECTIVENESS?

With Ertron,<sup>®</sup> local and systemic improvement has been reported in 701 out of 852 arthritic patients. Thus beneficial results were obtained in 82%; and no improvement was noted in only 17.8%. In a disease as resistant as rheumatoid arthritis, this is truly effective therapy.<sup>1</sup>

"No specificity is claimed for Ertron therapy. However, any substance which is non-toxic and which by its general systemic action does produce improved sense of well-being, diminution of soft tissue swelling, relief of pain, and improved muscle strength and which does make possible a return to gainful occupation, should be used in the treatment of arthritic patients."<sup>2</sup>


# ERTRON

STERIOD COMPLEX, WHITTIER

## TOLERABILITY?

Tolerance to Ertron is high in patients under periodic observation. Untoward side reactions are rare.<sup>3</sup> In 1,020 arthritic patients, marked intolerance requiring cessation of therapy occurred in only 1.4%, while minor side effects, such as nausea, gastrointestinal upset, headache, etc., were encountered in about 8%.

"These mild digestive disturbances disappear almost immediately after the cessation of Ertron administration and usually do not recur when this therapy is again instituted."<sup>4</sup>



Ertron is supplied in bottles of 50, 100 and 500 capsules, and Ertron Parenteral in packages of six 1 cc. ampuls. Each capsule contains 5 milligrams of activation-products having antirachitic activity of fifty thousand U.S.P. units. Each ampul contains activation-products having antirachitic activity of five hundred thousand U.S.P. units, in sesame oil. Biologically standardized.

## BIBLIOGRAPHY

- (1) Magnuson, P. B.; McElvenney, R. T., and Logan, E. E.: J. Michigan M. Soc. 46:71, 1947
- (2) Levinthal, D. H.; Logan, C. E.; Kohn, K. H., and Fishbein, W. I.: Indust. Med. 13:337, 1944
- (3) Cohen, A., and Reinhold, J. G.: Indust. Med. 17:442, 1948
- (4) Farley, R. T.; Spierling, H. F., and Kraines, S. H.: Indust. Med. 10:341, 1941

*Whittier*

LABORATORIES • DIVISION NUTRITION RESEARCH LABORATORIES, CHICAGO 30, ILLINOIS






## What Would You Say?

Twice a month we will select a caption for this cartoon from those sent in by our readers and send the author \$5. This caption was written by

Calvin B. Kitchen,  
M.D.  
Cleveland

Mail your caption to  
The Cartoon Editor,  
MODERN MEDICINE,  
84 South 10th St.,  
Minneapolis 3, Minn.



"Well, you know you can deduct the full amount from your income tax."




## Lewis Chair-Table

For General Examination  
and Treatment

- Hydraulic hand-pump elevation.
- Positioning from chair to table in one movement.
- Comfortably padded plastic upholstery.
- Porcelain enamel base—chrome plated trim.
- Adjustable head and foot rests.

◀ Chair  
Position

Table ▶  
Position



Send coupon for details

Name..... Street.....

City..... State..... Dealer's Name.....

**SHAMPAIN CO.** - 1926 SOUTH JEFFERSON  
ST. LOUIS 4, MISSOURI

*the fluid sulfadiazine that's*

*...better tasting*

*...faster acting*



To many patients—children, the aged and those with sore throats—swallowing bulky, half-gram sulfadiazine tablets is one of the discomforts of being sick. But these patients take ESKADIAZINE willingly. It tastes good.

It is not thick and cloying; it is light and easy to swallow.

Each 5 cc. (one teaspoonful) of ESKADIAZINE contains 0.5 Gm. (7.7 gr.) sulfadiazine—the dosage equivalent of the standard sulfadiazine tablet.

Yet desired serum levels are attained 3 to 5 times more rapidly with ESKADIAZINE than with sulfadiazine tablets. This is ascribed to the fact that ESKADIAZINE contains sulfadiazine in microcrystalline form.

*Smith, Kline & French Laboratories, Philadelphia*

***Eskadiazine***

*the outstandingly palatable fluid sulfadiazine*

# Current Books & Pamphlets

*This catalogue is computed from all available sources, American and foreign, to insure a complete listing of the month's releases.*

## Medicine

AN EPIHOME OF THE LABORATORY DIAGNOSIS AND TREATMENT OF TROPICAL DISEASES by Horace M. Shelley. 2d ed. 147 pp. Staples Press, New York City. \$2.25

DIFFERENTIAL DIAGNOSIS OF CHEST DISEASES by Jacob Jesse Singer. 344 pp., ill. Lea & Febiger, Philadelphia. \$7.50

## Surgery

AN ATLAS OF AMPUTATIONS by Donald B. Slocum. 562 pp., ill. C. V. Mosby Co., St. Louis. \$20

LES PROBLÈMES DU TRAITEMENT DES ULCÈRES PERFORÉS GASTRODUODÉNAUX: LA GASTRECTOMIE D'URGENCE-INDICATIONS ET RÉSULTATS by J. Mialaret and G. Edelmann. 136 pp. Masson & Co., Paris 320 fr.

## Psychiatry & Psychology

KORSCHACH'S TEST VOLUME I, BASIC PROCESSES by Samuel J. Beck. 2d ed. 242 pp. Grune & Stratton, New York City. \$4  
MONGOLISM AND CRETINISM by Clemens E. Benda. 2d ed. 333 pp., ill. Grune & Stratton, New York City. \$7



**TREAT ECZEMA  
THIS  
IMPROVED  
NEW WAY**

*Prescribe*

**SUPERTAH**

(Nason's)

*Crude Coal Tar Streamlined*

SUPERTAH Ointment is a white non-staining ointment prepared from a crude coal tar concentrate, uniformly milled in proper proportions to equal either a 5% or 10% crude tar ointment.

"It has proven as valuable as the black coal tar preparation and the advantage of the diminution of the black color is perfectly obvious."\*

\*Swartz & Reilly, "Diagnosis and Treatment of Skin Diseases", p. 66.

**TAILBY-NASON COMPANY • BOSTON 42, MASS.**



Today, as in 1934  
When symptoms indicate  
**Cow's Milk  
Lactalbumin  
Allergy**  
prescribe  
Meyenberg Evaporated  
Goat Milk

Rx 1950

It's Natural Milk


It's Nutritionally Equivalent to  
Evaporated Cow's Milk  
May be used in all formulae and recipes designating  
evaporated cow's milk - with no change in caloric  
value, fat or protein content.

Its Taste Approximates  
Evaporated Cow's Milk

Of particular importance in the patient beyond the  
infant formula stage.

It's Prepared Under Exact Control  
From milking to packing, constant control assures  
uniformity, purity. The milk is sterilized after sealing  
in the cans.

Available all year 'round at most drug stores in 14  
oz. cans. Advertised to the profession only.



By the makers of Hi-Pro® High-Protein  
Low-Fat Powdered Cow's Milk.



For further information, literature, for-  
mula feeding cards, write  
**Special Milk Products, Inc.**  
Los Angeles 64, California • Since 1934

DEVOTED TO RESEARCH AND CLINICAL  
DISEASES AND PROCESSES OF THE AGED

# Geriatrics

*Official Journal of the*  
**AMERICAN GERIATRICS SOCIETY**

Join the thousands of your  
colleagues who read  
**GERIATRICS** for the most  
significant developments in re-  
search and clinical study of the  
diseases of the aged and aging.  
**GERIATRICS** keeps you abreast  
of progress in diagnosis and  
treatment of cardiovascular,  
neurological, arteriosclerotic,  
arthritic, proctologic, pros-  
tatic, nutritive and other  
disorders.

**Subscribe NOW! — Save \$5 on BARGAIN  
INTRODUCTORY OFFER — 3 YEARS, \$10!**

**GERIATRICS**, 84 South 10th St., Minneapolis 3, Minn.

I'd like to have **GERIATRICS** help me keep abreast of  
latest developments in geriatrics. I'll take your bargain sub-  
scription deal.

Name.....Address.....

City.....Zone.....State.....

Check Enclosed..... Please Bill Me.....

**GERIATRICS** is  
chosen by the  
most eminent  
authors to pre-  
sent their im-  
portant papers  
on geriatrics.

### Gynecology

ESSENTIALS OF GYNECOLOGY by Leo Brady *et al.* 2d ed. 256 pp., ill. Macmillan Co., New York City. \$3

### Anesthesia

MODERN PRACTICE IN ANAESTHESIA edited by Frankis T. Evans. 566 pp., ill. Butterworth & Co., London. 50s.

### Cardiovascular Diseases

ULCERS AND ECZEMA OF THE LEG, SEQUELS OF PHLEBITIS, ETC: STUDIES ON STATIC DISEASES OF THE LOWER LIMBS AND THEIR TREATMENT by Holger Ove Bisgaard. 112 pp., ill. Ejnar Munksgaard, Copenhagen. 15 kr.

DISEASES OF THE HEART by Charles K. Friedberg. 1,081 pp., ill. W. B. Saunders Co., Philadelphia. \$11.50

DISEASES OF THE AORTA: DIAGNOSIS AND TREATMENT by Nathaniel F. Reich. 288 pp., ill. Macmillan Co., New York City. \$7.50

### Pediatrics

THE PSYCHOANALYTIC STUDY OF THE CHILD, 1947-1948 edited by Phyllis Greenacre *et al.* 493 pp. International Universities Press, New York City. \$10

THE RETARDED CHILD by Herta Loewy. 52 pp. Staples Press, London. 3s. 6d.

HAEMOLYTIC DISEASES OF THE NEWBORN by M. M. Pickles. 192 pp., ill. Blackwell Scientific Publications, Oxford. 21s.

### Allied Sciences

PHYSIKALISCHE CHEMIE IN MEDIZIN UND BIOLOGIE by W. Bladergroen. 2d ed. 675 pp., ill. B. Wepf & Co., Basel, Switzerland. 45 Sw. fr.


JORDAN-BURROWS' TEXTBOOK OF BACTERIOLOGY by William Burrows *et al.* 15th ed. 981 pp., ill. W. B. Saunders Co., Philadelphia. \$9

THE ELEMENTS OF GENETICS by Cyril Dean Darlington and Kenneth Mather. 446 pp. Macmillan Co., New York City. \$3.75

## Gebauer's S.S. SPRAY

### NEW SOAPLESS DETERGENT

- NON-IRRITATING
- AUTOMATICALLY DISPENSED



Gebauer's S. S. Spray provides a thick, creamy lather without water or waste. It is indicated for use in the following applications:

1. For preparing the field of operation, prior to use of antiseptics in minor surgery.
2. For the removal of ointment dressings.
3. As a cleansing agent for soap-sensitive patients.
4. For the rapid removal of oil and grease from the injuries of industrial patients.

S. S. Spray can be used instantly. It is packaged in Dispenseal bottle that prevents contamination when not in use. Now available at your local surgical supply dealers.

## THE GEBAUER CHEMICAL COMPANY

9410 St. Catherine Ave.

CLEVELAND, OHIO



**The NEWEST COMBINATION!**  
**MEDICS Ford-Bowles**  
**STETHOSCOPE**



No. P610  
**\$7.50**

**FORD BOWLES**  
 Valve control at  
 Your finger tip

At your Surgical Dealer  
 or Write for Literature  
**GRAHAM-FIELD**  
 273 Pearl St., N. Y. 7

## SLOW COLLECTIONS

Speed them up with the MODERN COLLECTION SYSTEM. Dr. A. L. Brown writes: "Used your SYSTEM just 30 days and it paid for itself several times." Write today for our FREE TRIAL OFFER.

Creditors Assn., 1477 Ridgeway Rd., Dayton 9, Ohio

**EVEREST & JENNINGS**  
**FOLDING WHEEL**  
**CHAIRS**  
**LIGHTEST and STRONGEST**



**FOLDS  
 TO 10  
 INCHES**



Everest & Jennings folding Wheel Chairs are **LIGHTEST AND STRONGEST** of all! They fold compactly for travel, work, play. Beautifully designed of chromium plated tubular steel. Your dealer can supply you, or write.

Manufacturers of the new  
 revolutionary WING FOLDING  
 ALUMINUM CRUTCHES

**EVEREST & JENNINGS, Dept. 17**

761 North Highland Ave.  
 LOS ANGELES 38, CALIF.

### Urology

UROLOGICAL ASPECTS OF SPINAL CORD INJURIES by Georg C. Prather. 146 pp., ill. Charles C. Thomas, Springfield, Ill. \$3.75

### Biography

MEMORIES OF EIGHTY YEARS by James Bryan Herrick. 270 pp., ill. University of Chicago Press, Chicago. \$5  
 A TWENTIETH CENTURY PHYSICIAN by Sir Arthur Hurst. 208 pp., ill. Edward Arnold & Co., London. 15s.

### Aviation Medicine

AVIATION MEDICINE: ITS THEORY AND APPLICATION by Kenneth G. Bergin. 448 pp., ill. John Wright & Sons, Bristol, England. 35s.

### Lay Reading

CHILDBIRTH: YOUR QUESTIONS ANSWERED by Carl Henry Davis and Donita Ferguson. 183 pp. Harper & Brothers, New York City. \$2.50  
 YOUR WEIGHT AND HOW TO CONTROL IT edited by Morris Fishbein. 276 pp. Doubleday & Co., Garden City, N.Y. \$2.95

### Public Health

HEALTH EDUCATION IN SCHOOLS by Jesse Feiring Williams and Ruth Abernathy. 316 pp. Ronald Press, New York City. \$3.50

### Nursing

RED CROSS NURSE IN ACTION 1882-1948 by Portia B. Kernodle. 524 pp. Harper & Bros., New York City. \$5  
 MOSBY'S COMPREHENSIVE REVIEW OF NURSING edited by a panel of specialists. 704 pp., ill. C. V. Mosby Co., St. Louis. \$5.75  
 HANDBOOK OF SURGICAL UROLOGY FOR INTERNS, HOSPITAL CORPSMEN AND NURSES by Nelse F. Ockerblad. 189 pp. Williams & Wilkins Co., Baltimore. \$3

### Miscellaneous

TONICS AND SEDATIVES: SELECTIONS FROM HIS FAVORITES edited by Morris Fishbein. 120 pp., ill. J. B. Lippincott Co., Philadelphia. \$2  
 THE ART OF READABLE WRITING by Rudolf F. Flesch. 237 pp. Harper & Brothers, New York City. \$3

Only on your  
Rx...

When prescribing Ergoapiol (Smith) with Savin for your gynecologic patients, you have the assurance that it can be obtained only on a written prescription, since this is the only manner in which this ethical preparation can be legally dispensed by the pharmacist. The dispensing of this **uterine tonic**, time-tested ERGOAPIOL (Smith) WITH SAVIN—only on your prescription—serves the best interests of physician and patient.

**INDICATIONS:** Amenorrhea, Dysmenorrhea, Menorrhagia, Metrorrhagia, and to aid involution of the postpartum uterus

**GENERAL DOSAGE:** One to two capsules, three to four times daily—as indications warrant.

In ethical packages of 20 capsules each, bearing no directions

## ERGOAPIOL (SMITH) WITH SAVIN

Literature Available  
to Physicians Only.



Ethical protective mark, M.H.S.,  
visible only when capsule  
is cut in half at seam.

MARTIN H. SMITH COMPANY  
150 Lafayette Street - New York 13, N. Y.



## FOR SALT RESTRICTED DIETS



Cellu Unsalted Vegetables—ready to heat and serve. All popular varieties. Write for the Cellu Catalog.

## CELLU Purposefully Prepared Dietary Foods

CHICAGO DIETETIC SUPPLY HOUSE, Inc.  
1750 West Van Buren Street Chicago 12, Ill.

## ARTHRITIS



One Gelucap Weapon For 2-Way Therapy

Many cases of arthritis are accompanied by fibrositis. Steinberg showed 143 out of 145 cases of primary fibrositis aided by high potency vitamin E. EDREX offers both vitamin D and vitamin E. ("An. Int. Med.", 19:136-139) Send for free literature.

**EDREX** VITAMINE E  
PLUS  
VITAMIN D  
**WILCO LABORATORIES**  
800 N. Clark St., Chicago 10, Ill.



Your women patients will welcome the prompt relief from pain and distress when you prescribe

## HAYDEN'S VIBURNUM COMPOUND



Professional  
Samples  
on Request

Excellent as a sedative, non-toxic and non-laxative, HVC produces the desired results without the unpleasant after effects caused by hypnotics. Relieves smooth muscle spasms. Safe for intestinal cramps.

**NEW YORK PHARMACEUTICAL COMPANY**  
Bedford Springs Bedford, Mass.

## PATIENTS

### ... I Have Met

The editors will pay \$1 for each story published. No contributions will be returned. Send your experiences to the Patients I Have Met Editor, MODERN MEDICINE, 84 South Tenth St., Minneapolis 3, Minn.

### The Full Circle

I was once a patient at a famous clinic. I was so grateful for all they did for me that I felt I owed the head physician something more than his fee, so I sent him an expensive engraved fountain pen for Christmas.

A year or so later I was passing through his town, so stopped in his office just to say "Hello." While there I had occasion to sign some papers and found that there was no ink in my pen. The famous surgeon handed me his. It was the pen I had sent to him. I was proud to think that he still cherished and used it.

As I screwed the cap back on and started to hand it back, he said, "You might need it later. Just take it along with you. I've got a whole drawer full that I'll never use. Patients have sent them to me as presents. That's one I haven't used."

I pocketed it and went on my way.  
—P. H.



"Your whole phobia seems to revolve about an unearthly fear of wringer wash machines."

*Announcing...*

**'PERAZIL'<sup>1</sup>**  
BRAND

CHLORCYCLIZINE HYDROCHLORIDE

*a (new type) antihistaminic*

*Relief from allergic symptoms for  
12 to 24 hours with a single dose*



'Perazil' brand Chlorcyclizine Hydrochloride is a completely new type of antihistaminic, its distinctive component being a piperazine ring instead of the usual ethylenediamine grouping. This uniquely different chemical structure results in a prolongation of action—up to 24 hours following a single 50 mg. dose.<sup>1</sup> In contrast to many other antihistaminic compounds, 'Perazil' exhibits a low incidence of side-effects despite its high potency and prolonged effectiveness.

**INDICATIONS:** Hay fever, vasomotor rhinitis, urticaria, allergic dermatitis and pollen asthma.

**DOSAGE:** 50 mg. (one product) once daily with water; may be increased to two or three times daily if required in very severe cases.

**PREPARATION:** 'Perazil' brand Chlorcyclizine Hydrochloride 50 mg. Compressed (scored). Bottles of 100.

<sup>1</sup> Jaros, S. H.: *Annals of Allergy*, Vol. 7, No. 4 (July-Aug.) 1949



**BURROUGHS WELLCOME & CO.** (U. S. A.) INC., TUCKAHOE 7, N. Y.

## CHLORTUSSIN (ANGIER)

a valuable adjunct treatment for coughs due to common cold, bronchitis, pharyngitis, laryngitis, pertussis. Especially designed for children. Contains B. Bottles of 4 and 8 fl. ozs.

ANGIER CHEMICAL CO., BOSTON 34, MASS.

Makers of

**THI-BO-MIN** — pleasant, non-alcoholic iron-liver preparation plus B<sub>1</sub> for supplemental treatment of Deficiency Anemias. Suitable for children.

**T-BARDRIN** — rectal suppositories for quick and prolonged symptomatic relief in Asthma—Hay Fever.



## AR-EX FOUNDATION LOTION

### 2-WAY AID FOR SKIN BLEMISHES...

**PSYCHOLOGICAL**—Hides minor blemishes of acne.

**MEDICAL**—Provides an excellent vehicle for sulfur, resorcinol, salicylic acid, etc.

**3 COSMETIC SHADES**

Send for professional sample

1036-M W. Van Buren St.

Ar-Ex Cosmetics, Inc. Chicago 7, ILL.

## UROPATHY

R Relief  
R Sedation  
R Bacteriostasis



Formula / Fluid oz.  
Methenamine . . . 18 gr.  
Sandalwood . . . 30 gr.  
Saw Palmetto . . . 30 gr.  
Zea . . . . . 30 gr.

Alcohol 9%

Available on prescription only, in 8-oz. bottles.

FREE  
SAMPLE

DRUG SPECIALTIES, INC.

210 Boyd Street, Los Angeles 54, Calif.

Professional Sample, Please:

M. D.

## In a Tough Spot

A friend of mine who practices in Miami Beach was complaining that his medical practice was twice as difficult as that of physicians located in other parts of the country.

"Just think," he exclaimed, "I can never tell my patients to go to Florida for their health."—M.C.




"I never realized these shots would be such a pain in the er-ah-ahem . . ."

## Porty Line

The following incident occurred in my practice recently.

The woman's voice on the country telephone sounded faint as she asked what to do for her father, who was constipated. It was impossible to carry on a conversation because the line kept fading, and our voices would cease to be heard. My last words to her were "Give him an enema," and then I futilely hung up, afraid that the woman had not heard my directions.

A few minutes later, the phone rang again. I couldn't distinguish what the woman was saying, but I was taking no chances this time. I shouted, "Give him an enema." Imagine my dismay when the voice, now suddenly clear and obviously that of another woman, reported acidly, "That's the first time I was told to give an enema for an ingrown toenail!"—A.S.



## *In neuromuscular dysfunction*

Physotropin is an important adjunct in the treatment of neuromuscular dysfunction, as it tends to facilitate nerve impulse transmission.

Physotropin employs the antagonism between Physostigmine and Atropine to remove the undesirable actions of the former without restricting its effect on the cranial nerves and skeletal muscles. Prescribe Physotropin. Your pharmacist can supply it.

**Indications:** Rheumatoid Arthritis • Bursitis • Anterior Poliomyelitis • Traumatic Neuromuscular Dysfunction • Myasthenia Gravis

**Supplied:** *Injectable Solution of Physotropin is supplied in 10 cc Rub-R-Top vials and Physotropin tablets in containers of 100, 500 and 1,000.*

▼ Write for professional samples and literature.

# physotropin

DURST

• S. F. DURST & CO., INC. • PHILADELPHIA 20, PENNA.





Sample on Request

**DOMECHEMICALS, INC.** 109 W. 64 ST.  
N. Y. 23, N. Y.



## LITTLE TOIDEY for Training the Baby

Little Toidey, \$5.50, and Toddler's Toidey (base and pan) \$3.95, are most convenient in the office of physicians who have patients with babies. 20% COURTESY DISCOUNT to doctors on all orders mailed directly to us. Write for complete list and free book "Training the Baby."

## THE TOIDEY COMPANY

Gertrude A. Muller, Inc.  
FORT WAYNE • INDIANA

## TUBULIN

Trade Mark

DEPROTEINATED RENAL EXTRACT

## VASODILATOR

Useful in treatment of:

### Peripheral Vascular Diseases

1. Arteriosclerosis Obliterans.
2. Buerger's Disease.
3. Raynaud's Syndrome.
4. Vaso-spasm.

Dosage: One c. c. Every Four Days.  
Subcutaneously and Intramuscularly.  
Supplied in Ampoules: 1.0 c. c.

Each 1 c. c. represents sterile aqueous solution of deproteinated renal extract obtained from 2.5 grams of Kidney tissue. For Literature write

## INCRETO PRODUCTS CORP.

231 W. 58th STREET NEW YORK

## INDEX TO ADVERTISERS

The publishers are not responsible for any errors or omissions.

Abbott Laboratories	107
American Cystoscope Makers, Inc.	99
American Ferment Company, Inc.	79
American Sterilizer Company	28
Angier Chemical Co.	126
Ar-Ex Cosmetics, Inc.	126
Arlington Chemical Company, The	36-37
Barnes, A. C., Company	11
Rauer & Black	81
Belmont Laboratories	30
Birtcher Corporation, The	40
Borden Company, The	66-67
Burroughs Wellcome & Co. (U. S. A.)	125
Central Pharmaceutical Co., The	83
Chatham Pharmaceuticals, Inc.	27
Chicago Dietetic Supply House, Inc.	124
Chilcott Labs. Div. of Maltine Co.	5
Ciba Pharmaceutical Products, Inc.	23, 4th Cover
Creditors Association	122
Dome Chemicals, Inc.	128
Drug Specialties, Inc.	126
Durst, S. F., & Co., Inc.	127
Eaton Laboratories, Inc.	75
Everest & Jennings	122
Florida Citrus Commission	111
Gebauer Chemical Company, The	121
Gerber Products Co.	113
Geriatrics	129
Graham-Ffield	122
Hoffman-La Roche, Inc.	8
Increto Products Corp.	128
Johnson & Johnson	86-87
Knox Gelatine	78
Kremers-Urban Company	92-93
Lederle Laboratories, Inc.	33
Leitz, E., Inc.	101
McNeil Laboratories, Inc.	3
Merck & Co., Inc.	19
Merrell, The Wm. S., Company	2nd Cover
M & R Dietetic Laboratories, Inc.	105
National Drug Company, The	29, 130
National Synthetics, Inc.	77
New York Pharmaceutical Company	124
Petra Manufacturing Co.	112
Professional Drugs, Inc.	4
Professional Printing Company, Inc.	31
Robins, A. H., Company, Inc.	15, 85, 102-103
Rorrig, J. B., & Company	89
Rystan Co., Inc.	43
Schenley Laboratories, Inc.	21
Schleffelin & Co.	107
Seck & Kade, Inc.	35
Shampaine Company	116
Sharp & Dohme	12-13
Sherman Laboratories	95
Shield Laboratories	22-23
Smith, Kline & French Laboratories	7, 34, 91, 117
Smith, Martin H., Company	125
Special Milk Products, Inc.	119
Talby-Nason Company	118
Toidey Company, The	128
Ulmer Laboratories	129
U. S. Vitamin Corporation	41
Valsey-Bristol Shoe Co., Inc.	32
Varick Pharmaceutical Co., Inc.	104
Viobin Laboratories	39
Walker Vitamin Products, Inc.	17
Wampole, Henry K., & Co., Inc.	42
White Laboratories, Inc.	18, 109
Whitehall Pharmaceutical Company	97
Whitely Laboratories	115
Wileo Laboratories	124
Winthrop-Stearns, Inc.	3rd Cover
Wyeth Incorporated	44

**FOR THE DOCTOR'S  
OFFICE PRACTICE**

**BILIPOSOL**  
TRADE MARK REG. U. S. PAT. OFF.

**A LIPOID SOLUBLE BASIC BISMUTH  
IN A CLEAR HOMOGENEOUS OIL SOLUTION**

**Safe Intramuscular Injection  
for the Treatment of**

**SYPHILIS**

**Immediate and Continued  
Dominance Over Syphilis**

**NO PAIN, TOXICITY OR SHOCK**

**NO HOSPITALIZATION REQUIRED**

**BILIPOSOL HAS STOOD THE TEST OF TIME**

**BILIPOSOL, in Boxes of 12, 50 and 100 ampoules, each containing 8 ctg. bismuth,  
IS OBTAINABLE FROM LEADING PHYSICIANS' SUPPLY  
HOUSES, RETAIL AND WHOLESALE DRUGGISTS OR FROM**

**LITERATURE MAILED TO PHYSICIANS ON REQUEST**

**ULMER LABORATORIES**

**414 So. Sixth Street Minneapolis, Minnesota**

**Sole General Distributors for United States and Canada**



**in arthritis**

• *unlike gold and other drugs*

**safe detoxifying colloidal sulfur**

• *relieves pain*

• *reduces joint swelling*

• *produces clinical remission  
even in the most severe cases*

**Sulphocol<sup>®</sup>** *colloidal sulfur compound*

**ORAL AND PARENTERAL**



Sulphocol Capsules (5 gr.) 1 or 2 after meals. Bottles of 100. Sulphocol Sol (parenteral), 25 cc. vials; 12 and 100-2 cc. vials. 1/4 to 1/2 cc. intramuscularly at 3 to 7 day intervals, gradually increased to 3 cc. Write for literature and samples of Sulphocol Capsules.

**A Product of the Mulford Colloid Laboratories  
THE NATIONAL DRUG COMPANY, PHILADELPHIA 44, PA.**



*Manufacturers of Pharmaceutical, Biological and Biochemical Products for the Medical Profession*

**SALT**

**without**

**SODIUM**



**NEOCURTASAL®**

When cardiac failure, hypertension, arteriosclerosis, or pregnancy complications call for a sodium free diet, you can let your patients have **salt without sodium: Neocurtasal**, the completely sodium free seasoning agent. Neocurtasal looks and is used like regular table salt.

Constituents: Potassium chloride, ammonium chloride, potassium formate, calcium formate, magnesium citrate and starch. Potassium content 36%; chloride 39.3%; calcium 0.3%; magnesium 0.2%.

Available in convenient  
2 oz. shakers and 8 oz. bottles.

NEOCURTASAL,  
trademark reg.  
U. S. & Canada


NEOCURTASAL,  
trademark reg. U. S. & Canada

*Winthrop Stearns* INC.  
NEW YORK 12, N. Y. WINDSOR, ONT.

# Pyribenzamine

**For control of allergic symptoms which may accompany a Common Cold . . .**

Pyribenzamine hydrochloride has been effectively and widely used in the control of allergic symptoms. Clinical tests have repeatedly shown that Pyribenzamine provides maximum allergic relief with minimal side effects. For relief of the allergic symptoms which may be associated with the common cold, Pyribenzamine is available in the following forms:



## **1. Pyribenzamine-Ephedrine for systemic treatment**

Each tablet contains 25 mg. of Pyribenzamine hydrochloride and 12 mg. of ephedrine sulfate. This combination synergistically promotes decongestion of the entire respiratory tract including the nasopharyngeal mucosa.



## **2. Pyribenzamine Nebulizer to control nasal symptoms**

Immediate relief with no systemic side effects. Pocket-size nebulizer distributes a mist of minute droplets of Pyribenzamine hydrochloride Nasal Solution 0.5% throughout the nasal passages.



## **3. Pyribenzamine Expectorant to control cough**

Each teaspoonful contains 30 mg. of Pyribenzamine citrate, 10 mg. of ephedrine sulfate and 80 mg. of ammonium chloride. Highly effective for relief of cough. Blocks congestive and spasmogenic effects of histamine shrinks respiratory mucosa and liquefies bronchial secretions.

More prescriptions have been written for Pyribenzamine than for all other antihistaminics.

# Ciba

**PHARMACEUTICAL PRODUCTS, INC., SUMMIT, NEW JERSEY**

**PYRIBENZAMINE** (brand of tripeleannamine) Trade Mark Reg. U. S. Pat. Off. 2/16031

MODERN MEDICINE  
84 S. 10 St. Minneapolis 3, Minn.

POSTMASTER: If undeliverable FOR ANY REASON  
notify sender, stating reason on form 3547.

Sec. 34.66, P. L. & R.  
U. S. POSTAGE  
**PAID**  
PERMIT No. 29  
Long Prairie, Minn.

WASHINGTON 25 D C  
HQARS U S AIR FORCE  
DR JACK BOLLERUD  
1219A